

1264942

THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE

United States Patent and Trademark Office

December 21, 2004

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE.

APPLICATION NUMBER: 60/518,166

FILING DATE: *November 07, 2003*

RELATED PCT APPLICATION NUMBER: PCT/US04/37242

Certified by



Jon W Dudas

Acting Under Secretary of Commerce
for Intellectual Property
and Acting Director of the U.S.
Patent and Trademark Office

PROVISIONAL APPLICATION TRANSMITTAL LETTER	Docket No. 3492	Type a plus sign (+) inside this box →
---	-----------------	---

To the Commissioner for Patents:

This is a request for filing a **PROVISIONAL APPLICATION** under 37 CFR 1.53(c).

Inventor(s)/Applicant(s)			
Last Name	First Name	Initial	Residence Address (City & either State or Foreign Country)
Zhou	Hongxing		Bellevue, Washington
Carter	Paul		Seattle, Washington

Title of the Invention (280 characters max):
ANTIBODIES THAT BIND INTERLEUKIN-4 RECEPTOR

Enclosed application parts are:

- ☒ 68 pages of specification, 10 pages of claim(s), 1 pages of abstract, and
pages of sequence listing, totaling 79 pages
- ☒ 8 sheets of drawings Figures 1-3
- ☐ Other (Specify) _____
- ☐ Other (Specify) _____

Method of Payment:

- ☒ Provisional filing fee amount \$160.00
- ☐ Check or money order enclosed to cover Provisional filing fee(s)
- ☒ Please charge Deposit Account No. 09-0089, in the name of Immunex Corporation, in the amount of \$160.00.
An original and one copy are enclosed.
- ☒ The Commissioner is hereby authorized to charge any additional fees which may be required or credit any
overpayment to Deposit Account No. 09-0089.

Respectfully submitted,



Nathan A. Machin
Attorney/Agent for Applicant(s)
Registration No.: 47,763
Phone: (206) 265-6779
Date: November 7, 2003

Please send all future correspondence to:

Immunex Corporation
Law Department
51 University Street
Seattle, Washington 98101
(206) 587-0430



22932

PATENT TRADEMARK OFFICE

22582 U.S. PTO
60/518166
110703

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In the Application of:
Hongxing Zhou and Paul Carter

Attorney Docket No. : 3492

Serial No.: --to be assigned--

Filed: November 7, 2003

For: ANTIBODIES THAT BIND INTERLEUKIN-4 RECEPTOR

CERTIFICATE OF MAILING BY EXPRESS MAIL

MAIL STOP PROVISIONAL PATENT APPLICATION

Commissioner for Patents

P.O. Box 1450

Alexandria, VA 22313-1450

EXPRESS MAIL LABEL NUMBER: EV 352011877 US

I hereby certify that the enclosed correspondence is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 C.F.R. §1.10 on the date listed below, and is addressed to MAIL STOP PROVISIONAL PATENT APPLICATION, Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.

Provisional Application for Patent Cover Sheet (+ copy)
Specification, Claims and Abstract (79 pages)
Drawings (8 sheets, Figures 1-3)
Postcard

Signed: _____


Kathleen F. Prindle

Date: _____

11/7/03

5 **TITLE**

Antibodies That Bind Interleukin-4 Receptor

BACKGROUND OF THE INVENTION

Interleukin-4 (IL-4), previously known as B cell stimulating factor, or BSF-1, was originally
 10 characterized by its ability to stimulate the proliferation of B cells in response to low concentrations
 of antibodies directed to surface immunoglobulin. IL-4 has been shown to possess a far broader
 spectrum of biological activities, including growth co-stimulation of T cells, mast cells,
 granulocytes, megakaryocytes, and erythrocytes. In addition, IL-4 stimulates the proliferation of
 15 several IL-2- and IL-3-dependent cell lines, induces the expression of class II major
 histocompatibility complex molecules on resting B cells, and enhances the secretion of IgE and
 IgG1 isotypes by stimulated B cells. IL-4 is associated with a TH2-type immune response, being
 one of the cytokines secreted by TH2 cells.

Murine and human IL-4 have been identified and characterized, including cloning of IL-4
 cDNAs and determination of the nucleotide and encoded amino acid sequences. *See Yokota et*
 20 *al.*, 1986, *Proc. Natl. Acad. Sci. USA* 83:5894; *Noma et al.*, 1986, *Nature* 319:640; *Grabstein et al.*,
 1986, *J. Exp. Med.* 163:1405; and U.S. Patent 5,017,691.

IL-4 binds to particular cell surface receptors, which results in transduction of a biological
 signal to cells such as various immune effector cells. IL-4 receptors are described, and DNA and
 amino acid sequence information presented, in *Mosley et al.*, 1989, *Cell* 59:335-48, (murine IL-
 25 4R); *Idzerda et al.*, 1990, *J. Exp. Med.* 171:861-73, (human IL-4R); and U.S. Patent 5,599,905.
 The IL-4 receptor described in these publications is sometimes referred to as IL-4R alpha.

Other proteins have been reported to be associated with IL-4R alpha on some cell types,
 and to be components of multi-subunit IL-4 receptor complexes. One such subunit is IL-2R
 gamma, also known as IL-2R gamma c. *See the discussion of IL-4R complexes in Sato et al.*,
 30 1994, *Current Opinion in Cell Biology*, 6:174-79. IL-4R alpha also has been reported to be a
 component of certain multi-subunit IL-13 receptor complexes. *See Zurawski et al.*, 1995, *J. Biol.*
Chem. 270:13869; *de Vries*, 1998, *J. Allergy Clin. Immunol.* 102:165; and *Callard et al.*, 1996,
Immunology Today, 17:108.

IL-4 has been implicated in a number of disorders, examples of which are allergy and
 35 asthma.

SUMMARY OF THE INVENTION

The present invention provides methods and compositions relating to novel IL-4R
 antagonists, in particular antibodies and antibody derivatives, that bind IL-4R alpha.

40 In one aspect, the present invention provides an antibody, wherein the light chain variable
 domain comprises a sequence of amino acids that differs from SEQ ID NO:4 only by: at least one
 amino acid substitution selected from the group consisting of S28T, S30N, S30G, S31N, S32D,
 S32N, A52T, S54Y, T57P, T57S, G93D, S94H, S94R, P96A, P97G, and T99M, and, optionally,
 one or more amino acid substitutions selected from the group consisting of E1D, L4M, S7T, G9A,
 45 K40R, F50Y, S68F, S77T, V86I, K105R, V106L, and E107D, and/or the heavy chain variable

5 domain comprises a sequence of amino acids that differs from SEQ ID NO:16 only by: at least one amino acid substitution selected from the group consisting of N58S, Y101W, F102Y, D103T, D103N, D103P, Y104H, Y104N, Y104W, and Y104R, and, optionally, one or more amino acid substitutions selected from the group consisting of Q6E, H13Q, G24A, R86S, and M90T, wherein said antibody binds to IL-4 receptor alpha. In one embodiment, the light chain CDR1 comprises a
 10 sequence selected from the group consisting of residues 24-35 of SEQ ID NO:6, residues 24-35 of SEQ ID NO:8, residues 24-35 of SEQ ID NO:10, residues 24-35 of SEQ ID NO:12; and residues 24-35 of SEQ ID NO:14, the light chain CDR2 comprises a sequence selected from the group consisting of residues 51-57 of SEQ ID NO:4, residues 51-57 of SEQ ID NO:6, residues 51-57 of SEQ ID NO:10, and residues 51-57 of SEQ ID NO:12, the light chain CDR3 comprises a
 15 sequence selected from the group consisting of residues 90-99 of SEQ ID NO:4, residues 90-99 of SEQ ID NO:6, residues 90-99 of SEQ ID NO:8, and residues 90-99 of SEQ ID NO:14, the heavy chain CDR1 comprises the sequence residues 31-35 of SEQ ID NO:16, the heavy chain CDR2 comprises a sequence selected from the group consisting of residues 50-65 of SEQ ID NO:16, and residues 50-65 of SEQ ID NO:18, and/or the heavy chain CDR3 comprises a
 20 sequence selected from the group consisting of residues 98-104 of SEQ ID NO:16, residues 98-104 of SEQ ID NO:18, residues 98-104 of SEQ ID NO:20, residues 98-104 of SEQ ID NO:22, residues 98-104 of SEQ ID NO:24, residues 98-104 of SEQ ID NO:26, residues 98-104 of SEQ ID NO:30, and residues 98-104 of SEQ ID NO:34. In another embodiment, the light chain FR1 comprises a sequence selected from the group consisting of residues 1-23 of SEQ ID NO:4,
 25 residues 1-23 of SEQ ID NO:10, residues 1-23 of SEQ ID NO:12, and residues 1-23 of SEQ ID NO:14, the light chain FR2 comprises a sequence selected from the group consisting of residues 36-50 of SEQ ID NO:4, and residues 36-50 of SEQ ID NO:14, the light chain FR3 comprises a sequence selected from the group consisting of residues 58-89 of SEQ ID NO:4, residues 58-89 of SEQ ID NO:10, and residues 58-89 of SEQ ID NO:12, the light chain FR4 comprises a
 30 sequence selected from the group consisting of residues 100-109 of SEQ ID NO:4, residues 100-109 of SEQ ID NO:8, and residues 100-109 of SEQ ID NO:12, the heavy chain FR1 comprises a sequence selected from the group consisting of residues 1-30 of SEQ ID NO:16, and residues 1-30 of SEQ ID NO:42, the heavy chain FR2 comprises the sequence of residues 1-30 of SEQ ID NO:16, the heavy chain FR3 comprises a sequence selected from the group consisting of
 35 residues 66-97 of SEQ ID NO:16, residues 66-97 of SEQ ID NO:18, and residues 66-97 of SEQ ID NO:42, and/or the heavy chain FR4 comprises the sequence of residues 105-115 of SEQ ID NO:16.

In another aspect, the present invention provides an antibody comprising: a light chain variable domain comprising a sequence that is at least 80% identical to a sequence selected from
 40 the group consisting of SEQ ID NO:4, 6, 8, 10, 12, and 14, with the proviso that said light chain variable domain does not comprise the sequence of SEQ ID NO:4, and/or a heavy chain variable domain comprising a sequence that is at least 80% identical to a sequence selected from the group consisting of SEQ ID NO:16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, and 62, with the proviso that said heavy chain variable domain does not
 45 comprise the sequence of SEQ ID NO:16, wherein said antibody binds to IL-4 receptor alpha. In

5 one embodiment, said light chain variable domain comprises a sequence that is at least 85% identical to a sequence selected from the group consisting of SEQ ID NO:4, 6, 8, 10, 12, and 14, and/or said heavy chain variable domain comprises a sequence that is at least 85% identical to a sequence selected from the group consisting of SEQ ID NO:16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, and 62. In another embodiment, said light chain variable domain comprises a sequence that is at least 90% identical to a sequence selected from the group consisting of SEQ ID NO:4, 6, 8, 10, 12, and 14, and/or said heavy chain variable domain comprises a sequence that is at least 90% identical to a sequence selected from the group consisting of SEQ ID NO:16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, and 62. In another embodiment, said light chain variable domain comprises a sequence that is at least 95% identical to a sequence selected from the group consisting of SEQ ID NO:4, 6, 8, 10, 12, and 14, and/or said heavy chain variable domain comprises a sequence that is at least 95% identical to a sequence selected from the group consisting of SEQ ID NO:16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, and 62. In another embodiment, said light chain variable domain comprises a sequence that is selected from the group consisting of SEQ ID NO:6, 8, 10, 12, and 14, and/or said heavy chain variable domain comprises a sequence that is selected from the group consisting of SEQ ID NO:16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, and 62.

In another aspect, the present invention provides an antibody wherein the light chain variable domain comprises a sequence of at least 15 contiguous amino acids of a sequence selected from the group consisting of: SEQ ID NO:4, 6, 8, 10, 12, and 14, with the proviso that said light chain variable domain does not comprise the sequence of SEQ ID NO:4, and/or wherein the heavy chain variable domain comprises a sequence of at least 15 contiguous amino acids of a sequence selected from the group consisting of SEQ ID NO:16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, and 62, with the proviso that said heavy chain variable domain does not comprise the sequence of SEQ ID NO:16, wherein said antibody binds to IL-4 receptor alpha. In one embodiment, the light chain variable domain comprises a sequence of at least 20 contiguous amino acids of a sequence selected from the group consisting of: SEQ ID NO:4, 6, 8, 10, 12, and 14, and/or the heavy chain variable domain comprises a sequence of at least 20 contiguous amino acids of a sequence selected from the group consisting of SEQ ID NO:16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, and 62. In another embodiment, the light chain variable domain comprises a sequence of at least 25 contiguous amino acids of a sequence selected from the group consisting of: SEQ ID NO:4, 6, 8, 10, 12, and 14, and/or the heavy chain variable domain comprises a sequence of at least 25 contiguous amino acids of a sequence selected from the group consisting of: SEQ ID NO:16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, and 62. In another embodiment, the light chain variable domain comprises a sequence of at least 35 contiguous amino acids of a sequence selected from the group consisting of: SEQ ID NO:4, 6, 8, 10, 12, and 14, and/or the heavy chain variable domain comprises a sequence of at least 35 contiguous amino acids of a sequence selected from the group consisting of: SEQ ID NO:16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, and 62. In another embodiment, the

5 light chain variable domain comprises a sequence of at least 50 contiguous amino acids of a sequence selected from the group consisting of: SEQ ID NO:4, 6, 8, 10, 12, and 14, and/or the heavy chain variable domain comprises a sequence of at least 50 contiguous amino acids of a sequence selected from the group consisting of: SEQ ID NO:16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, and 62. In another embodiment, the light chain
10 variable domain comprises a sequence of at least 75 contiguous amino acids of a sequence selected from the group consisting of: SEQ ID NO:4, 6, 8, 10, 12, and 14, and/or the heavy chain variable domain comprises a sequence of at least 75 contiguous amino acids of a sequence selected from the group consisting of: SEQ ID NO:16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, and 62. In another embodiment, the light chain variable
15 domain comprises a sequence of at least 100 contiguous amino acids of a sequence selected from the group consisting of: SEQ ID NO:4, 6, 8, 10, 12, and 14, and/or the heavy chain variable domain comprises a sequence of at least 100 contiguous amino acids of a sequence selected from the group consisting of: SEQ ID NO:16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, and 62.

20 In another aspect, the present invention provides an antibody comprising: a light chain variable domain comprising an amino acid sequence that is encoded by a nucleotide sequence that is at least 80% identical to a nucleotide sequence selected from the group consisting of SEQ ID NO:3, 5, 7, 9, 11, and 13, and/or a heavy chain variable domain comprising an amino acid sequence that is encoded by a nucleotide sequence that is at least 80% identical to a nucleotide
25 sequence selected from the group consisting of SEQ ID NO:15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59, and 61, with the proviso that said antibody does not comprise both the light chain variable domain sequence of SEQ ID NO:4 and the heavy chain variable domain sequence of SEQ ID NO:16, and wherein said antibody binds to IL-4 receptor alpha. In another embodiment, said light chain variable domain comprises an amino acid
30 sequence that is encoded by a nucleotide sequence that is at least 85% identical to a nucleotide sequence selected from the group consisting of SEQ ID NO:3, 5, 7, 9, 11, and 13, and/or said heavy chain variable domain comprises an amino acid sequence that is encoded by a nucleotide sequence that is at least 85% identical to a nucleotide sequence selected from the group consisting of SEQ ID NO:15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51,
35 53, 55, 57, 59, and 61. In another embodiment, said light chain variable domain comprises an amino acid sequence that is encoded by a nucleotide sequence that is at least 90% identical to a nucleotide sequence selected from the group consisting of SEQ ID NO:3, 5, 7, 9, 11, and 13, and/or said heavy chain variable domain comprises an amino acid sequence that is encoded by a nucleotide sequence that is at least 90% identical to a nucleotide sequence selected from the
40 group consisting of SEQ ID NO:15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59, and 61. In another embodiment, said light chain variable domain comprises an amino acid sequence that is encoded by a nucleotide sequence that is at least 95% identical to a nucleotide sequence selected from the group consisting of SEQ ID NO:3, 5, 7, 9, 11, and 13, and/or said heavy chain variable domain comprises an amino acid sequence that is encoded by a
45 nucleotide sequence that is at least 95% identical to a nucleotide sequence selected from the

5 group consisting of SEQ ID NO:15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59, and 61. In another embodiment, said light chain variable domain comprises an amino acid sequence that is encoded by a nucleotide sequence that is at least 98% identical to a nucleotide sequence selected from the group consisting of SEQ ID NO:3, 5, 7, 9, 11, and 13, and/or said heavy chain variable domain comprises an amino acid sequence that is encoded by a
10 nucleotide sequence that is at least 98% identical to a nucleotide sequence selected from the group consisting of SEQ ID NO:15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59, and 61. In another embodiment, said light chain variable domain comprises an amino acid sequence that is encoded by a nucleotide sequence selected from the group consisting of SEQ ID NO:5, 7, 9, 11, and 13, and/or said heavy chain variable domain comprises
15 an amino acid sequence that is encoded by a nucleotide sequence selected from the group consisting of SEQ ID NO:16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, and 62.

In another aspect, the present invention provides an antibody comprising: a light chain variable domain comprising an amino acid sequence that is encoded by a nucleotide sequence
20 that hybridizes under moderately stringent conditions to the complement of a nucleotide sequence selected from the group consisting of SEQ ID NO:3, 5, 7, 9, 11, and 13, and/or a heavy chain variable domain comprising an amino acid sequence that is encoded by a nucleotide sequence that hybridizes under moderately stringent conditions to the complement of a nucleotide sequence selected from the group consisting of SEQ ID NO:15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39,
25 41, 43, 45, 47, 49, 51, 53, 55, 57, 59, and 61, with the proviso that said antibody does not comprise both the light chain variable domain sequence of SEQ ID NO:4 and the heavy chain variable domain sequence of SEQ ID NO:16, and wherein said antibody binds to IL-4 receptor alpha. In one embodiment, said light chain variable domain comprises an amino acid sequence that is encoded by a nucleotide sequence that hybridizes under stringent conditions to the
30 complement of a nucleotide sequence selected from the group consisting of SEQ ID NO:3, 5, 7, 9, 11, and 13, and/or said heavy chain variable domain comprises an amino acid sequence that is encoded by a nucleotide sequence that hybridizes under stringent conditions to the complement of a nucleotide sequence selected from the group consisting of SEQ ID NO:15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59, and 61.

35 In another aspect, the present invention provides an isolated antibody, wherein the light chain variable region of said antibody comprises an amino acid sequence selected from the group consisting of SEQ ID NO:6, 8, 10, 12, and 14, and said antibody binds to IL-4 receptor alpha.

In another aspect, the present invention provides an isolated antibody, wherein the heavy chain variable region of said antibody comprises an amino acid sequence selected from the group
40 consisting of SEQ ID NO:16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, and 62, and said antibody binds to IL-4 receptor alpha.

In another aspect, the present invention provides an antibody selected from the group consisting of L2H1, L3H1, L4H1, L5H1, L1H2, L1H3, L1H4, L1H5, L1H6, L1H7, L1H8, L1H9, L1H10, L1H11, L2H4, L2H12, L2H13, L2H14, L6H1, L2H2, L2H3, L2H6, L2H7, L2H8, L2H9,
45 L2H10, and L2H11.

5 In another aspect, the present invention provides a human, humanized, or chimeric antibody.

 In another aspect, the present invention provides a monoclonal antibody.

 In another aspect, the present invention provides an antibody selected from the group consisting of an IgD, IgE, IgM, IgG1, IgG2, IgG3, IgG4, and IgG4 having at least one mutation in a
10 hinge region that alleviates a tendency to form intra-H chain disulfide bond antibody.

 In another aspect, the present invention provides an isolated polypeptide comprising an IL-4 receptor binding portion of an antibody of the invention. In one embodiment, said polypeptide comprises a Fab, F(ab')₂, scFv, diabody, triabody, or tetrabody.

 In another aspect, the present invention provides an isolated nucleic acid comprising a
15 nucleotide sequence encoding the light chain of an antibody of the invention, the heavy chain of an antibody of the invention, or a polypeptide of the invention.

 In another aspect, the present invention provides a vector comprising an isolated nucleic acid of the invention. In another embodiment, said vector is an expression vector.

 In another aspect, the present invention provides an isolated cell comprising a nucleic acid
20 of the invention. In one embodiment, said cell is a hybridoma. In another embodiment, said cell is a transgenic cell.

 In another aspect, the present invention provides a cell comprising an antibody of the invention. In one embodiment, said cell is a hybridoma. In another embodiment, said cell is a transgenic cell.

25 In another aspect, the present invention provides a cell comprising a polypeptide of the invention. In one embodiment, said cell is a transgenic cell.

 In another aspect, the present invention provides a method of making an antibody of the invention comprising incubating a cell comprising a nucleic acid encoding the light chain of said antibody and a nucleic acid encoding the heavy chain of said antibody under conditions that allow
30 said cell to express said light chain and said heavy chain and that allow said light chain and said heavy chain to assemble into said antibody, and isolating said antibody from said cell. In one embodiment, said cell is a hybridoma. In another embodiment, said cell is a transgenic cell.

 In another aspect, the present invention provides a method of inhibiting an IL-4 receptor comprising contacting a cell expressing IL-4 receptor alpha with an antibody of the invention or a polypeptide of the invention under conditions that allow said antibody or said polypeptide to bind to
35 said IL-4 receptor alpha. In one embodiment, said cell is a human cell. In another embodiment, said human cell is in a human.

 In another aspect, the present invention provides a method of treating a condition in a subject comprising administering to said subject an amount of an antibody of the invention or of a polypeptide of the invention effective for treating said condition. In one embodiment, said
40 condition is an inflammatory or cancerous condition. In another embodiment, said inflammatory or cancerous condition is an immunological condition. In another embodiment, said condition is asthma, septic arthritis, dermatitis herpetiformis, chronic idiopathic urticaria, ulcerative colitis, scleroderma, hypertrophic scarring, Whipple's Disease, benign prostate hyperplasia, a lung
45 disorder in which IL-4 receptor plays a role, condition in which IL-4 receptor-mediated epithelial

5 barrier disruption plays a role, a disorder of the digestive system in which IL-4 receptor plays a
 role, an allergic reaction to a medication, Kawasaki disease, sickle cell disease, Churg-Strauss
 syndrome, Grave's disease, pre-eclampsia, Sjogren's syndrome, autoimmune lymphoproliferative
 syndrome, autoimmune hemolytic anemia, Barrett's esophagus, autoimmune uveitis, tuberculosis,
 cystic fibrosis, allergic bronchopulmonary mycosis, chronic obstructive pulmonary disease,
 10 bleomycin-induced pneumopathy and fibrosis, radiation-induced pulmonary fibrosis, pulmonary
 alveolar proteinosis, adult respiratory distress syndrome, sarcoidosis, hyper IgE syndrome,
 idiopathic hypereosinophil syndrome, an autoimmune blistering disease, pemphigus vulgaris,
 bullous pemphigoid, myasthenia gravis, chronic fatigue syndrome, or nephrosis.

In another aspect, the present invention provides a pharmaceutical composition
 15 comprising an antibody of the invention or a polypeptide of the invention and an excipient, diluent,
 or buffer.

BRIEF DESCRIPTION OF THE DRAWINGS

Figures 1A-C present the nucleotide sequence of the coding region of a human IL-4
 20 receptor alpha cDNA. The amino acid sequence encoded by the cDNA is presented as well. The
 cDNA clone was isolated from a cDNA library derived from a human T cell line T22. The encoded
 protein comprises (from N- to C-terminus) an N-terminal signal peptide, followed by an
 extracellular domain, a transmembrane region (underlined), and a cytoplasmic domain, as
 discussed further in PCT publication WO 01/92340 A3. The nucleotide sequence and amino acid
 25 sequences of Figures 1A to 1C also are presented in SEQ ID NO:1 and 2, respectively.

Figures 2A-2D present polynucleotide sequences encoding the light chain variable regions
 of L1 (SEQ ID NO:3), L2 (SEQ ID NO:5), L3, (SEQ ID NO:7), L4 (SEQ ID NO:9), L5 (SEQ ID
 NO:11), and L6 (SEQ ID NO:13), and polynucleotide sequences encoding the heavy chain
 variable regions of H1 (SEQ ID NO:15), H2 (SEQ ID NO:17), H3 (SEQ ID NO:19), H4 (SEQ ID
 30 NO:21), H5 (SEQ ID NO:23), H6 (SEQ ID NO:25), H7 (SEQ ID NO:27), H8 (SEQ ID NO:29), H9
 (SEQ ID NO:31), H10 (SEQ ID NO:33), H11 (SEQ ID NO:35), H12 (SEQ ID NO:37), H13 (SEQ ID
 NO:39), H14 (SEQ ID NO:41), H15 (SEQ ID NO:43), H16 (SEQ ID NO:45), H17 (SEQ ID NO:47),
 H18 (SEQ ID NO:49), H19 (SEQ ID NO:51), H20 (SEQ ID NO:53), H21 (SEQ ID NO:55), H22
 (SEQ ID NO:57), H23 (SEQ ID NO:59), and H24 (SEQ ID NO:61). The sequences are shown
 35 using one-letter nucleotide abbreviations. Sequences corresponding to CDR1, CDR2, and CDR3
 regions are shown in bold type for each sequence and underlined in L1 and H1. Sequences
 corresponding to FR1, FR2, FR3, and FR4 are shown in plain type.

Figure 3 presents the amino acid sequences of the light chain variable regions of L1 (SEQ
 ID NO:4), L2 (SEQ ID NO:6), L3, (SEQ ID NO:8), L4 (SEQ ID NO:10), L5 (SEQ ID NO:12), and L6
 40 (SEQ ID NO:14), and polynucleotide sequences encoding the heavy chain variable regions of H1
 (SEQ ID NO:16), H2 (SEQ ID NO:18), H3 (SEQ ID NO:20), H4 (SEQ ID NO:22), H5 (SEQ ID
 NO:24), H6 (SEQ ID NO:26), H7 (SEQ ID NO:28), H8 (SEQ ID NO:30), H9 (SEQ ID NO:32), H10
 (SEQ ID NO:34), H11 (SEQ ID NO:36), H12 (SEQ ID NO:38), H13 (SEQ ID NO:40), H14 (SEQ ID
 NO:42), H15 (SEQ ID NO:44), H16 (SEQ ID NO:46), H17 (SEQ ID NO:48), H18 (SEQ ID NO:50),
 45 H19 (SEQ ID NO:52), H20 (SEQ ID NO:54), H21 (SEQ ID NO:56), H22 (SEQ ID NO:58), H23

5 (SEQ ID NO:60), and H24 (SEQ ID NO:62). The sequences of the L1 and H1 variable regions are shown using one-letter amino acid abbreviations. Other light chain and heavy chain variable sequences are indicated by dashes at residues where they are identical to L1 or H1 and with the appropriate one-letter amino acid abbreviation where they differ from L1 or H1. Sequences corresponding to CDR1, CDR2, and CDR3 regions are shown in bold type for each sequence and underlined in L1 and H1. Sequences corresponding to FR1, FR2, FR3, and FR4 are shown in plain type.

DETAILED DESCRIPTION OF THE INVENTION

15 The present invention provides compositions and methods relating to anti-IL-4 receptor (IL-4R) antibodies, including methods for treating certain conditions mediated by IL-4R, and for inhibiting biological activities of interleukin-4 (IL-4) and interleukin-13 (IL-13) *in vivo*. Compositions of the invention include, for example, anti-IL-4R antibodies, polypeptides, polynucleotides, cells comprising or expressing antibodies, polypeptides or polynucleotides of the invention, and pharmaceutical compositions, examples of which are provided below.

20 Polynucleotide and polypeptide sequences are indicated using standard one- or three-letter abbreviations. Unless otherwise indicated, polypeptide sequences have their amino termini at the left and their carboxy termini at the right and single-stranded nucleic acid sequences, and the top strand of double-stranded nucleic acid sequences, have their 5' termini at the left and their 3' termini at the right. A particular polypeptide or polynucleotide sequence also can be described by explaining how it differs from a reference sequence. For example, the phrase "a polypeptide sequence that differs from SEQ ID NO:4 at S28T" describes a polypeptide sequence that is identical to SEQ ID NO:4 except that the serine residue at position 28 of SEQ ID NO:4 is replaced by a threonine residue.

30 Polynucleotide and polypeptide sequences of particular light and heavy chain variable regions are shown in Figures 2 and 3, respectively, where they are labeled, for example, L1 ("light chain variable region 1"), H1 ("heavy chain variable region 1"), etc. Antibodies comprising a light chain and heavy chain from Figure 3 are indicated by combining the name of the light chain and the name of the heavy chain variable regions. For example, "L4H7," indicates an antibody comprising the light chain variable sequence of L4 and the heavy chain variable sequence of H7.

35 "Light chain variable domain (or region)," "heavy chain variable domain (or region)," "CDR1, 2, and 3" and "FR1, 2, 3, and 4" are defined according to the scheme of Kabat *et al.* in *Sequences of Proteins of Immunological Interest*, 5th Ed., US Dept. of Health and Human Services, PHS, NIH, NIH Publication no. 91-3242, 1991.

40 The "percent identity" of two polynucleotide or two polypeptide sequences is determined by comparing the sequences using the GAP computer program (a part of the GCG Wisconsin Package, version 10.3 (Accelrys, San Diego, CA)) using its default parameters.

A biological molecule (*e.g.*, a polypeptide, antibody, or nucleic acid) is "isolated" or "substantially purified" if it is sufficiently free of other biological molecules, cell debris, and other substances to be used in standard laboratory protocols (*e.g.*, a binding or hybridization assay).

5 Methods of substantially purifying polypeptides, antibodies, and nucleic acids are well-known in the art.

The terms "peptide," "polypeptide" and "protein" are used interchangeably throughout and refer to a molecule comprising two or more amino acid residues joined to each other by peptide bonds.

10 The terms "polynucleotide," "oligonucleotide" and "nucleic acid" are used interchangeably throughout and include DNA molecules (e.g., cDNA or genomic DNA), RNA molecules (e.g., mRNA), analogs of the DNA or RNA generated using nucleotide analogs (e.g., peptide nucleic acids and non-naturally occurring nucleotide analogs), and hybrids thereof. The nucleic acid molecule can be single-stranded or double-stranded. In one embodiment, the nucleic acid
15 molecules of the invention comprise a contiguous open reading frame encoding an antibody, or a fragment, derivative, mutein, or variant thereof, of the invention.

Two single-stranded nucleic acid molecules are "the complement" of each other if their sequences can be aligned in an anti-parallel orientation such that every nucleotide in one sequence is opposite its complementary nucleotide in the other sequence, without the introduction
20 of gaps, and without unpaired nucleotides at the 5' or the 3' end of either sequence. A nucleic acid molecule that is "complementary" to a given nucleotide sequence is one that is sufficiently complementary to the given nucleotide sequence that it can hybridize under moderately stringent conditions to the given nucleotide sequence. Thus, a nucleic acid can be complementary to another nucleic acid without being its complement.

25 A "vector" is a nucleic acid that can be used to introduce another nucleic acid linked to it into a cell. One type of vector is a "plasmid," which refers to a linear or circular double stranded DNA molecule into which additional nucleic acid segments can be ligated. Another type of vector is a viral vector (e.g., replication defective retroviruses, adenoviruses and adeno-associated viruses), wherein additional DNA segments can be introduced into the viral genome. Certain
30 vectors are capable of autonomous replication in a host cell into which they are introduced (e.g., bacterial vectors comprising a bacterial origin of replication and episomal mammalian vectors). Other vectors (e.g., non-episomal mammalian vectors) are integrated into the genome of a host cell upon introduction into the host cell, and thereby are replicated along with the host genome. An "expression vector" is a type of vector that can direct the expression of a chosen
35 polynucleotide.

A nucleotide sequence is "operably linked" to a regulatory sequence if the regulatory sequence affects the expression (e.g., the level, timing, or location of expression) of the nucleotide sequence. A "regulatory sequence" is a nucleic acid that affects the expression (e.g., the level, timing, or location of expression) of a nucleic acid. The regulatory sequence can, for example,
40 exert its effects directly on the regulated nucleic acid, or through the action of one or more polypeptides (e.g., polypeptides that bind to the regulatory sequence and/or the nucleic acid). Examples of regulatory sequences include promoters, enhancers and other expression control elements (e.g., polyadenylation signals). Further examples of regulatory sequences are described in, for example, Goeddel, 1990, Gene Expression Technology: Methods in Enzymology 185,
45 Academic Press, San Diego, CA and Baron *et al.*, 1995, Nucleic Acids Res. 23:3605-06.

5 A "host cell" is a cell that can be used to express a nucleic acid, *e.g.*, a nucleic acid of the invention. A host cell can be a prokaryote, for example, *E. coli*, or it can be a eukaryote, for example, a single-celled eukaryote (*e.g.*, a yeast or other fungus), a plant cell (*e.g.*, a tobacco or tomato plant cell), an animal cell (*e.g.*, a human cell, a monkey cell, a hamster cell, a rat cell, a mouse cell, or an insect cell) or a hybridoma. Examples of host cells include the COS-7 line of
10 monkey kidney cells (ATCC CRL 1651) (see Gluzman *et al.*, 1981, Cell 23:175), L cells, C127 cells, 3T3 cells (ATCC CCL 163), Chinese hamster ovary (CHO) cells or their derivatives such as Veggie CHO and related cell lines which grow in serum-free media (see Rasmussen *et al.*, 1998, Cytotechnology 28:31) or CHO strain DX-B11, which is deficient in DHFR (see Urlaub *et al.*, 1980, Proc. Natl. Acad. Sci. USA 77:4216-20), HeLa cells, BHK (ATCC CRL 10) cell lines, the
15 CV1/EBNA cell line derived from the African green monkey kidney cell line CV1 (ATCC CCL 70) (see McMahon *et al.*, 1991, EMBO J. 10:2821), human embryonic kidney cells such as 293, 293 EBNA or MSR 293, human epidermal A431 cells, human Colo205 cells, other transformed primate cell lines, normal diploid cells, cell strains derived from in vitro culture of primary tissue, primary explants, HL-60, U937, HaK or Jurkat cells. Typically, a host cell is a cultured cell that can be
20 transformed or transfected with a polypeptide-encoding nucleic acid, which can then be expressed in the host cell. The phrase "recombinant host cell" can be used to denote a host cell that has been transformed or transfected with a nucleic acid to be expressed. A host cell also can be a cell that comprises the nucleic acid but does not express it at a desired level unless a regulatory sequence is introduced into the host cell such that it becomes operably linked with the nucleic
25 acid. It is understood that the term host cell refers not only to the particular subject cell but to the progeny or potential progeny of such a cell. Because certain modifications may occur in succeeding generations due to either mutation or environmental influences, such progeny may not, in fact, be identical to the parent cell, but are still included within the scope of the term as used herein.

30 A "chimeric antibody" is an antibody in which a portion of the heavy and/or light chain is identical with, homologous to, or derived from an antibody from a particular species or belonging to a particular antibody class or subclass, while the remainder of the chain(s) is/are identical with, homologous to, or derived from an antibody (-ies) from another species or belonging to another antibody class or subclass. Also included are fragments of such antibodies, that exhibit the
35 desired biological activity (*i.e.*, the ability to specifically bind IL-4 receptor). See, U.S. Patent No. 4,816,567 and Morrison, 1985, Science 229:1202-07.

A "CDR grafted antibody" is an antibody comprising one or more CDRs derived from an antibody of a particular species or isotype and the framework of another antibody of the same or different species or isotype.

40 A "multi-specific antibody" is an antibody that recognizes more than one epitope on one or more antigens. A subclass of this type of antibody is a "bi-specific antibody" which recognizes two distinct epitopes on the same or different antigens.

A "variant" of a polypeptide (*e.g.*, an antibody) comprises an amino acid sequence wherein one or more amino acid residues are inserted into, deleted from and/or substituted into

5 the amino acid sequence relative to another polypeptide sequence. Variants of the invention include fusion proteins.

A "derivative" of a polypeptide is a polypeptide (*e.g.*, an antibody) that has been chemically modified in some manner distinct from insertion, deletion, or substitution variants, *e.g.*, via conjugation to another chemical moiety. Unless otherwise indicated, the term "antibody" includes, 10 in addition to antibodies comprising two full-length heavy chains and two full-length light chains, derivatives, variants, fragments, and muteins thereof, examples of which are described below.

A molecule (*e.g.*, an antibody) "specifically binds IL-4 receptor" if it binds to IL-4 receptor, or a fragment thereof, with at least 10 times higher affinity than the molecule binds to a polypeptide unrelated to IL-4 receptor.

15 An "antigen binding domain" or "antigen binding region" is the portion of an antibody molecule which contains the amino acid residues (or other moieties) that interact with an antigen and confer on the antibody its specificity and affinity for the antigen.

An "epitope" is the portion of a molecule that is bound by an antibody. An epitope can comprise non-contiguous portions of the molecule (*e.g.*, in a polypeptide, amino acid residues that 20 are not contiguous to one another in the polypeptide's sequence but that, together in the context of the molecule, are bound by an antibody).

Indications

In one aspect, the present invention provides methods of treating, preventing, curing, 25 relieving, or ameliorating a disease, disorder, condition, or illness. Among the conditions to be treated in accordance with the present invention are asthma, septic/reactive arthritis, dermatitis herpetiformis, chronic idiopathic urticaria, scleroderma, hypertrophic scarring, Whipple's Disease, benign prostate hyperplasia, lung disorders in which IL-4 plays a role, conditions in which IL-4-induced epithelial barrier disruption plays a role, disorders of the digestive system in which IL-4 30 plays a role, including inflammatory bowel disease and other inflammatory conditions in the gastrointestinal tract, allergic reactions to medication, Kawasaki disease, sickle cell disease (including sickle cell crisis), Churg-Strauss syndrome, Grave's disease, pre-eclampsia, Sjogren's syndrome, autoimmune lymphoproliferative syndrome, autoimmune hemolytic anemia, Barrett's esophagus, autoimmune uveitis, tuberculosis, and nephrosis, as described in more detail below. 35 IL-4R antagonists also find use as adjuvants to allergy immunotherapy and as vaccine adjuvants.

Examples of antibodies suitable for treating these conditions are described below, and include, *e.g.*, antibodies that bind IL-4R and inhibit the binding of IL-4 thereto. Particularly useful antibodies also inhibit the binding of IL-13 to IL-13 receptor (IL-13R). Particular embodiments of the invention include novel antibodies and antibody derivatives, fragments, muteins and variants, 40 polypeptides, nucleic acid molecules, cells, methods of making the preceding, methods of inhibiting IL-4R α , and methods of treating a subject, examples of which are described below.

5 In one aspect, the present invention provides methods comprising administering an anti-IL-4R α antibody to a subject. In one embodiment, the subject is afflicted with, or at risk for developing, a condition (including, *e.g.*, an illness, infection, injury, disease, or disorder) that is caused, induced, mediated, potentiated, exacerbated, or otherwise affected, directly or indirectly, by the activity of IL-4R α . Such conditions include, for example, conditions caused, induced,
10 mediated, potentiated, exacerbated, or otherwise affected, directly or indirectly, by IL-4 and/or IL-13. Other factors or cytokines also may play a role in such conditions.

The biological activities of IL-4 are mediated through binding to specific cell surface receptors, referred to as interleukin-4 receptors (IL-4R). IL-4-induced conditions include those arising from biological responses that result from the binding of IL-4 to a native IL-4 receptor on a
15 cell, or which may be inhibited or suppressed by preventing IL-4 from binding to an IL-4 receptor. Conditions that may be treated include, but are not limited to, medical disorders characterized by abnormal expression of IL-4 or of one or more components of an IL-4R or IL-13R (including, for example, overexpression, misexpression in a particular tissue or cell type, or misexpression at a particular developmental stage), or by an abnormal host response to IL-4 production. Further
20 examples are conditions in which IL-4-induced antibody production, or proliferation or influx of a particular cell type, plays a role. IL-4-induced disorders include those in which IL-4 induces upregulation of IL-4 receptors or enhanced production of another protein that plays a role in a disease (*e.g.*, another cytokine).

A method for treating a mammal, including a human subject, who has such a medical
25 disorder comprises administering an anti-IL-4R antibody, or derivative thereof, to the mammal or otherwise contacting an IL-4R of the mammal with the antibody or derivative, *e.g.*, in an *ex vivo* procedure. Conditions that may be treated in accordance with the present invention are described, for example, in USSN 09/847,816, filed May 1, 2001, the relevant disclosure of which is incorporated by reference herein. Such conditions include, but are not limited to, asthma,
30 septic/reactive arthritis, dermatitis herpetiformis, urticaria (especially chronic idiopathic urticaria), ulcers, gastric inflammation, mucosal inflammation, ulcerative colitis, Crohn's Disease, inflammatory bowel disease, other disorders of the digestive system in which IL-4 plays a role (*e.g.*, IL-4-induced inflammation of part of the gastrointestinal tract), conditions in which IL-4-induced barrier disruption plays a role (*e.g.*, conditions characterized by decreased epithelial
35 barrier function in the lung or gastrointestinal tract), scleroderma, hypertrophic scarring, Whipple's Disease, benign prostate hyperplasia, IL-4-induced pulmonary conditions (including those listed below), allergic reactions to medication, Kawasaki disease, sickle cell disease or crisis, Churg-Strauss syndrome, Grave's disease, pre-eclampsia, Sjogren's syndrome, autoimmune lymphoproliferative syndrome, autoimmune hemolytic anemia, Barrett's esophagus, autoimmune
40 uveitis, tuberculosis, nephrosis, pemphigus vulgaris or bullous pemphigoid (autoimmune blistering diseases), and myasthenia gravis (an autoimmune muscular disease).

Anti-IL-4R antibodies, and derivatives thereof, also find use as adjuvants to allergy immunotherapy and as vaccine adjuvants. Accordingly, an anti-IL-4R antibody may be employed as an adjuvant to allergy immunotherapy treatment. Anti-IL-4R antibody find further use as
45 vaccine adjuvants, such as adjuvants for cancer vaccines and infectious disease vaccines. The

- 5 use of IL-4 adjuvants, especially when directing the immune response toward a TH1 response would be beneficial in treating or preventing the disease in question.

Septic/reactive arthritis

10 An anti-IL-4R antibody may be employed in treating septic arthritis, which also is known as reactive arthritis or bacterial arthritis. Septic arthritis can be triggered by (result from, or develop subsequent to) infection with such microbes as *Staphylococcus aureus*, *Chlamydia trachomatis*, *Yersinia e.g.*, *Y. enterocolitica*, *Salmonella, e.g.*, *S. enteritidis*, *Shigella* and *Campylobacter*. *S. aureus* has been reported to be the major human pathogen in septic arthritis, responsible for the majority of cases.

15 IL-4 and IL-4-dependent Th2 responses play roles in promoting septic arthritis. Anti-IL-4R antibody can be employed in accordance with the invention to inhibit IL-4 and also to suppress the Th2 response in patients having septic arthritis or at risk for developing septic arthritis.

20 IL-4 increases bacterial burden and bacterial persistence in joints, by inhibiting clearance of the bacteria. Anti-IL-4R antibody may be employed to assist in the clearance of bacteria associated with reactive arthritis, thereby reducing clinical manifestations such as swelling in joints. Anti-IL-4R antibody may be administered to a human subject afflicted with septic arthritis, to reduce IL-4-mediated joint inflammation. In one approach, an antagonist is injected into a joint, *e.g.*, into synovial fluid in the knee.

25 The use of an anti-IL-4R antibody may benefit subjects having (or at risk for) septic arthritis by suppressing a TH2 response and promoting a TH1 response against the infection. TH2 cytokines may contribute to bacterial persistence in the joint, whereas a TH1 response plays a role in eliminating the bacteria.

30 The antibody may be administered to subjects infected with bacteria or other microbes such as those listed above, to prevent development of septic arthritis. An Antibody may be administered, for example, after diagnosis with such an infection, but before development of clinical symptoms of septic arthritis.

Whipple's Disease

35 *Tropheryma whippelii* is the causative bacterium for Whipple's Disease, also known as intestinal lipodystrophy and lipophagia granulomatosis. The disease is characterized by steatorrhea, frequently generalized lymphadenopathy, arthritis, fever, and cough. Also reported in Whipple's Disease patients are an abundance of "foamy" macrophages in the jejunal lamina propria, and lymph nodes containing periodic acid-schiff positive particles appearing bacilliform by electron microscopy (*Steadman's Medical Dictionary*, 26th Edition, Williams & Wilkins, Baltimore, MD, 1995).

40 The use of anti-IL-4R antibody may benefit subjects having (or at risk for developing) Whipple's Disease, by restoring a normal balance between the TH1 and TH2 components of the patient's immune response. Increased production of IL-4 (a TH2-type cytokine) and decreased levels of certain TH1-type cytokines have been associated with Whipple's Disease. TH2 cytokines
45 may contribute to bacterial persistence, whereas a TH1 response plays a role in clearing the

- 5 causative bacteria. IL-4R antagonists may be administered to subjects infected with *T. whippelii*, whether or not the subject exhibits clinical symptoms of Whipple's Disease.

Dermatitis herpetiformis

10 Dermatitis herpetiformis, also known as Duhring's disease, is a chronic skin condition characterized by blistering skin lesions, cutaneous IgA deposits, and itching. Patients have an immunobullous skin disorder with an associated gluten sensitive enteropathy, which is mediated by a Th2 immune response. Anti-IL-4R antibody is administered in accordance with the present invention, to inhibit IL-4 and the Th2 response, thus promoting healing of current lesions and reducing or preventing the formation of blisters on the extensor body surfaces.

15

Hypertrophic scarring

In accordance with the present invention, anti-IL-4R antibody is administered to subjects who have, or are susceptible to developing, hypertrophic scarring. In one method provided herein, an anti-IL-4R antibody is administered to a subject with a burn injury. An immune response to burns and other injury is believed to play a role in the pathogenesis of hypertrophic scarring. Increased production of TH2-type cytokines, including IL-4, and reduced levels of certain TH1-type cytokines have been reported in burn patients who have hypertrophic scarring. The use of anti-IL-4R antibodies may benefit subjects having (or at risk for developing) hypertrophic scarring, by suppressing a TH2-type immune response.

25

Urticaria

Urticaria, especially chronic forms thereof such as chronic idiopathic urticaria (CIU), may be treated with an anti-IL-4R antibody in accordance with the present invention. CIU patients have higher serum levels of IL-4 than controls, and may have a predominantly TH2-type cytokine profile. Mast cells and Th2-type T cells are implicated as primary effector cells in chronic urticaria. IL-4 stimulates mast cell proliferation. Mast cell degranulation leads to histamine release, subsequent erythema, eosinophilia, redness of skin, and itching. Anti-IL-4R antibodies are administered to inhibit IL-4 and reduce the TH2-type response, thereby helping to control a subject's urticaria.

35

Ulcerative colitis; other disorders of the gastrointestinal tract

IL-4 is implicated in the pathogenesis of ulcerative colitis. Th2-type cytokines including IL-4 may predominate in the colonic mucosa of patients with this disorder. The use of anti-IL-4R antibodies to suppress the TH2 response may alleviate this condition.

40

In addition to ulcerative colitis, other disorders of the gastrointestinal tract or digestive system may be treated with anti-IL-4R antibodies. Examples of such disorders include, but are not limited to, inflammatory bowel disease (IBD), with ulcerative colitis and Crohn's Disease being forms of IBD, gastritis, ulcers, and mucosal inflammation.

Any gastrointestinal condition in which IL-4 plays a role may be treated with an anti-IL-4R antibody in accordance with the present invention. For example, conditions involving IL-4-induced

45

5 inflammation of part of the gastrointestinal tract may be treated with an anti-IL-4R antibody. Particular embodiments are directed to treatment of chronic inflammatory conditions in the gastrointestinal tract.

Other embodiments are directed to conditions in which IL-4-induced barrier disruption plays a role, e.g., conditions characterized by decreased epithelial barrier function in at least a
10 portion of the gastrointestinal tract. Such conditions may, for example, involve damage to the epithelium that is induced by IL-4, directly or indirectly.

The intestinal epithelium forms a relatively impermeable barrier between the lumen and the submucosa. Disruption of the epithelial barrier has been associated with conditions such as inflammatory bowel disease. See the discussion in Youakim, A. and M. Ahdieh (*Am. J. Physiol.*
15 276 (*Gastrointest. Liver Physiol.* 39):G1279-G1288, 1999), hereby incorporated by reference in its entirety. A damaged or "leaky" barrier can allow antigens to cross the barrier, which in turn elicits an immune response that may cause further damage to gastrointestinal tissue. Such an immune response may include recruitment of neutrophils or T cells, for example. An anti-IL-4R antibody may be administered to inhibit undesirable stimulation of an immune response.

20

Lung disorders

Methods for treating IL-4-induced pulmonary disorders are provided herein. Such disorders include, but are not limited to, lung fibrosis, including chronic fibrotic lung disease, other conditions characterized by IL-4-induced fibroblast proliferation or collagen accumulation in the
25 lungs, pulmonary conditions in which a TH2-type immune response plays a role, conditions characterized by decreased barrier function in the lung (e.g., resulting from IL-4-induced damage to the epithelium), or conditions in which IL-4 plays a role in an inflammatory response (e.g., asthma).

Cystic fibrosis is characterized by the overproduction of mucus and development of
30 chronic infections. Inhibiting IL-4 and the Th2 response will reduce mucus production and help control infections such as allergic bronchopulmonary aspergillosis (ABPA).

Allergic bronchopulmonary mycosis occurs primarily in patients with cystic fibrosis or asthma, where a Th2 immune response is dominant. Inhibiting IL-4 and the Th2 response will help clear and control these infections.

35 Chronic obstructive pulmonary disease is associated with mucus hypersecretion and fibrosis. Inhibiting IL-4 and the Th2 response will reduce the production of mucus and the development of fibrosis thereby improving respiratory function and delaying disease progression.

Bleomycin-induced pneumopathy and fibrosis, and radiation-induced pulmonary fibrosis are disorders characterized by fibrosis of the lung which is manifested by the influx of Th2, CD4⁺
40 cells and macrophages, which produce IL-4 which in turn mediates the development of fibrosis. Inhibiting IL-4 and the Th2 response will reduce or prevent the development of these disorders.

Pulmonary alveolar proteinosis is characterized by the disruption of surfactant clearance. IL-4 increases surfactant product. Use of anti-IL-4R antibody will decrease surfactant production and decrease the need for whole lung lavage.

5 Adult respiratory distress syndrome (ARDS) may be attributable to a number of factors,
one of which is exposure to toxic chemicals. One patient population susceptible to ARDS is
critically ill patients who go on ventilators. ARDS is a frequent complication in such patients. Anti-
IL-4R antibody treatment may alleviate ARDS by reducing inflammation and adhesion molecules,
although methods for treating such subjects in accordance with the present invention are not
10 limited by a particular mechanism of action. Anti-IL-4R antibody may be used to prevent or treat
ARDS.

Sarcoidosis is characterized by granulomatus lesions. Use of anti-IL-4R antibody to treat
sarcoidosis, particularly pulmonary sarcoidosis, is contemplated herein.

15 Conditions in which IL-4-induced barrier disruption plays a role (e.g., conditions
characterized by decreased epithelial barrier function in the lung) may be treated with anti-IL-4R
antibody. Damage to the epithelial barrier in the lungs may be induced by IL-4 directly or indirectly.
The epithelium in the lung functions as a selective barrier that prevents contents of the lung lumen
from entering the submucosa. A damaged or "leaky" barrier allows antigens to cross the barrier,
which in turn elicits an immune response that may cause further damage to lung tissue. Such an
20 immune response may include recruitment of eosinophils or mast cells, for example. An anti-IL-
4R antibody may be administered to inhibit such undesirable stimulation of an immune response.

Anti-IL-4R antibodies may be employed to promote healing of lung epithelium, thus
restoring barrier function. Anti-IL-4R antibody may be employed to promote healing of lung
epithelium in asthmatics, for example. Alternatively, the antagonist is administered for
25 prophylactic purposes, to prevent IL-4-induced damage to lung epithelium.

Tuberculosis

A TH2-type immune response is implicated in playing a role in causing tissue damage
(e.g., necrosis of lung tissue) in tuberculosis (TB) patients. Elevated levels of IL-4 are associated
30 with TB. IL-4 production may be particularly elevated in cavitary tuberculosis (i.e., in TB patients
who have developed pulmonary cavities, which can be detected/visualized by such techniques as
radiographs of the chest).

Anti-IL-4R antibodies may benefit TB patients (especially those with cavitary TB) by
suppressing a TH2-type immune response. Methods for treating such subjects in accordance with
35 the present invention are not limited by a particular mechanism of action, however. Anti-IL-4R
antibody advantageously are administered in an amount that restores the desired balance
between the TH1 and TH2 components of the immune response, and reduces IL-4-induced tissue
damage in a patient.

Churg-Strauss syndrome

40 Churg-Strauss syndrome, a disease also known as allergic granulomatous angiitis, is
characterized by inflammation of the blood vessels in persons with a history of asthma or allergy,
and by eosinophilia. Anti-IL-4R antibodies may be administered to alleviate inflammation in
subjects with this syndrome. The use of anti-IL-4R antibodies to suppress a TH2-type immune
45 response, and to combat eosinophilia, would benefit the subjects.

5

Pre-eclampsia

Pre-eclampsia is a toxemia of late pregnancy. The condition is characterized by a sharp rise in blood pressure, generally accompanied by edema and albuminuria, during the third term of pregnancy.

10

Elevated TH1-type and TH2-type immune responses may play a role in the condition.

One method provided herein comprises administering an anti-IL-4R antibody to a pregnant woman who has developed pre-eclampsia. The anti-IL-4R antibody is administered in an amount, and for a period of time, sufficient to reduce the level of IL-4 (or of TH2-type cytokines collectively) to a level that is considered normal during pregnancy. In general, the anti-IL-4R antibody is

15

administered repeatedly throughout the duration of the pregnancy.

Scleroderma

Anti-IL-4R antibodies are administered to subjects with scleroderma in accordance with the invention. The antibodies reduce IL-4-induced collagen synthesis by fibroblasts in the patients.

20

The antibodies may be employed in preventing or reducing fibrosis in skin and lung tissues, as well as other tissues in which fibrosis occurs in scleroderma patients, suppressing collagen synthesis in such tissues, and in treating scleroderma-related pulmonary disease.

Benign Prostate Hyperplasia

25

Benign prostate hyperplasia (BPH), also known as benign prostate hypertrophy, may be treated with anti-IL-4R antibodies. While not wishing to be bound by a particular mechanism of action, administration of an anti-IL-4R antibody may benefit a subject with BPH by suppressing IL-4-induced inflammation, or by suppressing a TH2-type immune response.

30

Grave's Disease

Antibodies directed against thyrotropin receptor play an important role in Grave's Disease, a disorder characterized by hyperthyroidism. Studies of cytokine production in Grave's Disease patients show a shift toward a TH2-type cytokine response. Use of an anti-IL-4R antibody to suppress the TH2-type immune response, and suppress antibody production, would benefit

35

Grave's Disease patients.

Sickle Cell Disease

Sickle cell disease patients typically experience intermittent periods of acute exacerbation called crises, with the crises being categorized as anemic or vaso-occlusive. Anti-IL-4R antibodies find use in treating or preventing sickle cell crisis, especially in subjects with elevated IL-4 levels or in whom the immune response has shifted toward a TH2-type response. Sickle cell disease (especially sickle cell crisis) has been associated with increased susceptibility to infectious diseases, including bacterial infections. Administering anti-IL-4R antibodies to sickle cell disease patients may help the patient mount an immune response against infectious diseases.

45

5 Sjogren's syndrome

The autoimmune disease known as Sjogren's syndrome or sicca syndrome typically combines dry eyes and dry mouth with a disorder of the connective tissues, such as rheumatoid arthritis, lupus, scleroderma, or polymyositis. The vast majority of patients are middle age (or older) females. Sjogren's syndrome is an inflammatory disease of glands (*e.g.*, lacrimal and salivary glands) and other tissues of the body. The syndrome typically is associated with autoantibody production.

Anti-IL-4R antibodies may be administered to reduce the inflammatory response (such as inflammation of glands, including lacrimal glands) in such subjects. Anti-IL-4R antibodies may benefit Sjogren's syndrome patients by suppressing a TH2-type immune response. Methods for treating subjects in accordance with the present invention are not limited by a particular mechanism of action, however.

Autoimmune lymphoproliferative syndrome

Manifestations of autoimmune lymphoproliferative syndrome include lymphoproliferation and autoantibody production. Patients with the syndrome reportedly have an inherited deficiency in apoptosis. Anti-IL-4R antibodies may benefit subjects with this syndrome by suppressing a TH2-type immune response. Methods for treating such subjects in accordance with the present invention are not limited by a particular mechanism of action, however.

25 Autoimmune hemolytic anemia

Excessive IL-4 secretion, and a deficiency in TH1-type cytokines, are implicated in contributing to the pathogenesis of autoimmune hemolytic anemia. Anti-IL-4R antibodies are administered in accordance with the present invention, to benefit the patients by reducing autoantibody production, and by restoring a more normal balance between the TH1 and TH2 components of the immune response.

Autoimmune uveitis

Uveitis involves inflammation of the uvea (generally considered to include the iris, ciliary body, and choroid, considered together). Excess IL-4 secretion is implicated as playing a role in pathogenesis of this sight-threatening inflammatory eye disease. In accordance with the present invention, anti-IL-4R antibodies are administered to a subject with, or at risk for developing, uveitis. In one embodiment, anti-IL-4R antibodies are administered to an individual who has autoimmune uveoretinitis.

40 Kawasaki Disease

Also known as the mucocutaneous lymph node syndrome, Kawasaki disease (KD) mainly afflicts young children. The disease is characterized by particular changes in the mucus membranes lining the lips and mouth, and by enlarged, tender lymph glands. Symptoms typically include fever, conjunctivitis, inflammation of the lips and mucous membranes of the mouth, swollen glands in the neck, and a rash covering the hands and feet, leading to hardened, swollen

- 5 and peeling skin on hands and feet. In children with Kawasaki Disease (KD), inflammation of arteries (vasculitis) may develop. Due to the effect of the disease on the vascular system, KD reportedly is the main cause of acquired heart disease in children.

Anti-IL-4R antibodies may be administered to subjects with Kawasaki Disease. Excessive IL-4 secretion and a deficiency in TH1-type cytokines contribute to the pathogenesis of the
10 disease.

Barrett's esophagus

Barrett's esophagus is a condition characterized by alteration (subsequent to irritation) of the cells in the epithelial tissue that lines the lower portion of the esophagus. Frequent reflux of
15 the stomach contents into the esophagus, over time, can lead to Barrett esophagus. Patients with Barrett esophagus are at risk for developing esophageal cancer (e.g., adenocarcinoma). While not wishing to be bound by a particular mechanism of action, administration of an anti-IL-4R antibody may benefit a subject with Barrett's esophagus by suppressing a TH2-type immune response. In one embodiment, an anti-IL-4R antibody is administered to a subject with
20 esophagitis, to inhibit progression to Barrett's esophagus.

Nephrosis

Nephrosis, also known as nephrotic syndrome, is a kidney disease that is non-inflammatory and non-malignant. In the condition known as minimal change nephrosis,
25 glomerular damage (believed to arise from structural changes in glomerular visceral epithelial cells) results in abnormalities that include proteinuria. A TH2-type immune response (especially secretion of the TH2-type cytokines IL-4 and IL-13) are implicated as playing a role in pathogenesis of minimal change nephrosis.

Other indications

Additional examples of conditions that may be treated in accordance with the present invention include but are not limited to the following. Anti-IL-4R antibodies may be employed in treating or preventing hyper IgE syndrome, idiopathic hypereosinophil syndrome, allergic reactions to medication, autoimmune blistering diseases (e.g., pemphigus vulgaris or bullous pemphigoid),
35 myasthenia gravis (an autoimmune muscular disease), and chronic fatigue syndrome. Anti-IL-4R antibodies may be employed in treating GVHD; particular methods for treating GVHD in combination with other therapeutic agents are described below. Anti-IL-4R antibodies also find use in treating or preventing hepatotoxicity induced by drugs such as diclofenac (a non-steroidal anti-inflammatory drug).

40 An anti-IL-4R antibody may be employed as an adjuvant to allergy immunotherapy treatment. Anti-IL-4R antibodies find further use as vaccine adjuvants, such as adjuvants for cancer vaccines and infectious disease vaccines. The use of anti-IL-4R antibodies is especially advantageous when favoring a TH1-type immune response would be beneficial in preventing or treating the condition for which the vaccine is being administered. Anti-IL-4R antibodies may be

- 5 employed when reducing an antibody-mediated immune response and/or promoting a T-cell-mediated immune response is desired.

Anti-IL-4R antibodies

10 In one aspect, the present invention provides antibodies, and fragments, derivatives, muteins, and variants thereof, which bind to IL-4 receptor alpha, *e.g.*, human IL-4 receptor alpha.

Anti-IL-4R antibodies that may be employed in accordance with the present invention include antibodies that inhibit a biological activity of IL-4. Examples of such biological activities include associating with another receptor component (*e.g.*, IL-2R gamma or IL-13R alpha), binding (either alone or as part of a multimeric receptor complex) a signaling molecule (*e.g.*, IL-4 or IL-13),
15 and transducing a signal in response to binding a signaling molecule.

Different anti-IL-4R antibodies may bind to different domains or epitopes of IL-4R or act by different mechanisms of action. Examples include but are not limited to antibodies that interfere with binding of IL-4 to IL-4R or that inhibit signal transduction. The site of action may be, for example, intracellular (*e.g.*, by interfering with an intracellular signaling cascade) or extracellular.
20 An antibody need not completely inhibit an IL-4 induced activity to find use in the present invention; rather, antibodies that reduce a particular activity of IL-4 are contemplated for use as well.

The above-presented discussions of particular mechanisms of action for anti-IL-4R antibodies in treating particular diseases are illustrative only, and the methods presented herein are not bound thereby. The mechanisms of action by which anti-IL-4R antibodies ameliorate
25 diseases are not limited to those discussed above.

An anti-IL-4R antibody may inhibit an IL-4-mediated influx of cells involved in an immune or inflammatory response. An antibody may act by, for example, reducing proliferation, activation, migration, influx, or accumulation of a particular cell type, or by inhibiting a biological response directly or indirectly attributable to a particular cell type. Examples of particular cell types are
30 fibroblasts, mast cells, and eosinophils.

As discussed above, some conditions may be treated by suppressing a TH2-type immune response. IL-4R is associated with a TH2 response, and is one of the cytokines secreted by T-helper cells of type 2 (TH2 cells). An anti-IL-4R antibody may be administered to reduce a TH2-type immune response. The IL-4R antibody may be said to reduce proliferation of TH2 cells, to
35 suppress a TH2 response, to shift the immune response toward a TH1 response, or to favor a TH1-type response. Antagonists of other TH2-type cytokine(s), such as IL-5, IL-10, or IL-13, may be additionally administered to subjects who have a disorder involving elevated levels of such cytokines. Techniques for measuring the amount of such cytokines in a subject, *e.g.*, in the subject's serum, are well known.

40 One embodiment of the invention is directed to a method for inhibiting IL-4-induced damage to epithelium, comprising administering an anti-IL-4R antibody to a subject who has, or is at risk of developing, a condition in which IL-4-mediated epithelial barrier disruption plays a role.

Particular embodiments of methods provided herein comprise administering an anti-IL-4R antibody to inhibit IL-4-induced damage to epithelium in the gastrointestinal tract or lung. Such
45 methods may be employed to prevent epithelial damage, or to restore epithelial barrier function

5 (i.e., promote repair or healing of the epithelium). The ability of an anti-IL-4R antibody to inhibit IL-4-induced damage to epithelium may be confirmed in any of a number of suitable assays, such as those described herein.

Any inflammation associated with (or subsequent to) an infection also may be treated with an anti-IL-4R antibody. The antibody may be administered to inhibit any IL-4-induced component
10 of an inflammatory response resulting from microbial infection in the gastrointestinal tract, for example.

Combinations of two or more antibodies or antibody derivatives, or of an antibody or antibody derivative and one or more other IL-4, IL-13, IL-4R and/or IL-13R antagonists (as described, for example, in US Patents 5,599,905, 5,840,869, 5,856,296, 5,767,065, 5,717,072,
15 6,391,581, 5,710,023, Idzerda *et al.*, 1990, J. Exp. Med. 171:861-73, and Mosley *et al.*, 1989, Cell 59:335-48, incorporated herein by reference in their entireties) may be employed in methods and compositions of the present invention.

Oligonucleotide-directed site-specific mutagenesis procedures can be employed to provide an altered gene having particular codons altered according to the substitution, deletion, or
20 insertion required. Examples of techniques for making such alterations are described in Walder *et al.*, 1986, Gene 42:133; Bauer *et al.* 1985, Gene 37:73; Craik, BioTechniques, January 1985, 12-19; Smith *et al.*, 1981, *Genetic Engineering: Principles and Methods*, Plenum Press; and U.S. Patent Nos. 4,518,584 and 4,737,462. These and other methods can be used to make, for example, derivatives of anti-IL-4R antibodies that have, for example, increased affinity, avidity, or
25 specificity for IL-4R as compared to the underivatized antibody.

Other derivatives of anti-IL-4R antibodies within the scope of this invention include covalent or aggregative conjugates of anti-IL-4R antibodies, or fragments thereof, with other proteins or polypeptides, such as by expression of recombinant fusion proteins comprising heterologous polypeptides fused to the N-terminus or C-terminus of an anti-IL-4R antibody
30 polypeptide. For example, the conjugated peptide may be a heterologous signal (or leader) polypeptide, e.g., the yeast alpha-factor leader, or a peptide such as an epitope tag. Anti-IL-4R antibody-containing fusion proteins can comprise peptides added to facilitate purification or identification of the anti-IL-4R antibody (e.g., poly-His). An anti-IL-4R antibody polypeptide also can be linked to the FLAG peptide Asp-Tyr-Lys-Asp-Asp-Asp-Lys (DYKDDDDK) (SEQ ID
35 NO:69) as described in Hopp *et al.*, *Bio/Technology* 6:1204, 1988, and U.S. Patent 5,011,912. The FLAG peptide is highly antigenic and provides an epitope reversibly bound by a specific monoclonal antibody (mAb), enabling rapid assay and facile purification of expressed recombinant protein. Reagents useful for preparing fusion proteins in which the FLAG peptide is fused to a given polypeptide are commercially available (Sigma, St. Louis, MO).

40 Oligomers that contain one or more anti-IL-4R antibody polypeptides may be employed as IL-4R antagonists. Oligomers may be in the form of covalently-linked or non-covalently-linked dimers, trimers, or higher oligomers. Oligomers comprising two or more anti-IL-4R antibody polypeptides are contemplated for use, with one example being a homodimer. Other oligomers include heterodimers, heterotrimers, *etc.*

5 One embodiment is directed to oligomers comprising multiple anti-IL-4R antibody polypeptides joined *via* covalent or non-covalent interactions between peptide moieties fused to the anti-IL-4R antibody polypeptides. Such peptides may be peptide linkers (spacers), or peptides that have the property of promoting oligomerization. Leucine zippers and certain polypeptides derived from antibodies are among the peptides that can promote oligomerization of IL-4R polypeptides attached thereto, as described in more detail below.

10 In particular embodiments, the oligomers comprise from two to four anti-IL-4R antibody polypeptides. The anti-IL-4R antibody moieties of the oligomer may be in any of the forms described above, *e.g.*, variants or fragments. Preferably, the oligomers comprise anti-IL-4R antibody polypeptides that have IL-4R binding activity.

15 In one embodiment, an oligomer is prepared using polypeptides derived from immunoglobulins. Preparation of fusion proteins comprising certain heterologous polypeptides fused to various portions of antibody-derived polypeptides (including the Fc domain) has been described, *e.g.*, by Ashkenazi *et al.*, 1991, PNAS USA 88:10535; Byrn *et al.*, 1990, Nature 344:677; and Hollenbaugh *et al.*, 1992 "Construction of Immunoglobulin Fusion Proteins", in
20 *Current Protocols in Immunology*, Suppl. 4, pages 10.19.1 - 10.19.11.

One embodiment of the present invention is directed to a dimer comprising two fusion proteins created by fusing an IL-4R binding fragment of an anti-IL-4R antibody to the Fc region of an antibody. The dimer can be made by, for example, inserting a gene fusion encoding the fusion protein into an appropriate expression vector, expressing the gene fusion in host cells transformed
25 with the recombinant expression vector, and allowing the expressed fusion protein to assemble much like antibody molecules, whereupon interchain disulfide bonds form between the Fc moieties to yield the dimer.

The term "Fc polypeptide" as used herein includes native and mutein forms of polypeptides derived from the Fc region of an antibody. Truncated forms of such polypeptides containing the hinge region that promotes dimerization are also included. Fusion proteins comprising Fc moieties (and oligomers formed therefrom) offer the advantage of facile purification by affinity chromatography over Protein A or Protein G columns.

One suitable Fc polypeptide, described in PCT application WO 93/10151 (hereby incorporated by reference), is a single chain polypeptide extending from the N-terminal hinge region to the native C-terminus of the Fc region of a human IgG1 antibody. Another useful Fc polypeptide is the Fc mutein described in U.S. Patent 5,457,035 and in Baum *et al.*, 1994, EMBO J. 13:3992-4001. The amino acid sequence of this mutein is identical to that of the native Fc sequence presented in WO 93/10151, except that amino acid 19 has been changed from Leu to Ala, amino acid 20 has been changed from Leu to Glu, and amino acid 22 has been changed from
35 Gly to Ala. The mutein exhibits reduced affinity for Fc receptors.

In other embodiments, the variable portion of the heavy and/or light chains of an anti-IL-4R antibody may be substituted for the variable portion of an antibody heavy and/or light chain.

Alternatively, the oligomer is a fusion protein comprising multiple anti-IL-4R antibody polypeptides, with or without peptide linkers (spacer peptides). Among the suitable peptide linkers
40 are those described in U.S. Patents 4,751,180 and 4,935,233.

5 Another method for preparing oligomeric anti-IL-4R antibody derivatives involves use of a leucine zipper. Leucine zipper domains are peptides that promote oligomerization of the proteins in which they are found. Leucine zippers were originally identified in several DNA-binding proteins (Landschulz *et al.*, 1988, Science 240:1759), and have since been found in a variety of different proteins. Among the known leucine zippers are naturally occurring peptides and derivatives
10 thereof that dimerize or trimerize. Examples of leucine zipper domains suitable for producing soluble oligomeric proteins are described in PCT application WO 94/10308, and the leucine zipper derived from lung surfactant protein D (SPD) described in Hoppe *et al.*, 1994, FEBS Letters 344:191, hereby incorporated by reference. The use of a modified leucine zipper that allows for stable trimerization of a heterologous protein fused thereto is described in Fanslow *et al.*, 1994,
15 Semin. Immunol. 6:267-78. In one approach, recombinant fusion proteins comprising an anti-IL-4R antibody fragment or derivative fused to a leucine zipper peptide are expressed in suitable host cells, and the soluble oligomeric anti-IL-4R antibody fragments or derivatives that form are recovered from the culture supernatant.

20 Anti-IL-4R antibody polypeptides and fusion proteins described herein may be prepared by any of a number of conventional techniques. For example, anti-IL-4R antibody polypeptides may be purified from cells that naturally express them, or they may be produced in recombinant expression systems, using any technique known in the art.

Any expression system known in the art can be used to make the recombinant polypeptides of the invention. In general, host cells are transformed with a recombinant
25 expression vector that comprises DNA encoding a desired anti-IL-4R antibody polypeptide. Among the host cells that may be employed are prokaryotes, yeast or higher eukaryotic cells. Prokaryotes include gram negative or gram positive organisms, for example *E. coli* or bacilli. Higher eukaryotic cells include insect cells and established cell lines of mammalian origin. Examples of suitable mammalian host cell lines include the COS-7 line of monkey kidney cells
30 (ATCC CRL 1651) (Gluzman *et al.*, 1981, Cell 23:175), L cells, 293 cells, C127 cells, 3T3 cells (ATCC CCL 163), Chinese hamster ovary (CHO) cells, HeLa cells, BHK (ATCC CRL 10) cell lines, and the CV1/EBNA cell line derived from the African green monkey kidney cell line CV1 (ATCC CCL 70) as described by McMahon *et al.*, 1991, EMBO J. 10: 2821. Appropriate cloning and expression vectors for use with bacterial, fungal, yeast, and mammalian cellular hosts are
35 described by Pouwels *et al.* (*Cloning Vectors: A Laboratory Manual*, Elsevier, New York, 1985).

The transformed cells are cultured under conditions that promote expression of the anti-IL-4R antibody polypeptide, and the polypeptide is recovered by conventional protein purification procedures. One such purification procedure includes the use of affinity chromatography, *e.g.*,
over a matrix having all or a portion (*e.g.*, the extracellular domain) of IL-4R bound thereto.

40 Polypeptides contemplated for use herein include substantially homogeneous recombinant mammalian anti-IL-4R antibody polypeptides substantially free of contaminating endogenous materials.

In one aspect, the present invention provides antibodies that interfere with the binding of IL-4 to an IL-4 receptor. Such antibodies, referred to herein as blocking antibodies, may be raised
45 against IL-4R, or a fragment, variant or derivative thereof, and screened in conventional assays for

5 the ability to interfere with binding of IL-4 to IL-4 receptors. Examples of suitable assays are assays that test the antibodies for the ability to inhibit binding of IL-4 to cells expressing IL-4R, or that test antibodies for the ability to reduce a biological or cellular response that results from the binding of IL-4 to cell surface IL-4 receptors.

10 It has been reported that IL-4R alpha is a component of certain multi-subunit IL-13 receptor complexes (Zurawski *et al.*, 1995, J. Biol. Chem. 270: 13869; de Vries, 1998, J. Allergy Clin. Immunol. 102:165; and Callard *et al.*, 1996, Immunology Today, 17:108, each incorporated by reference herein). Accordingly, in one embodiment, an antibody is provided that blocks binding of IL-4 and also of IL-13 to cells. The antibodies inhibit IL-4-induced biological activity and also inhibit IL-13-induced activity, and thus may be employed in treating conditions induced by
15 either or both cytokines. Examples of such conditions include but are not limited to IgE-mediated conditions, asthma, allergic conditions, allergic rhinitis, and dermatitis including atopic dermatitis.

Antibodies that bind to IL-4R alpha may be screened in various conventional assays to determine whether they interfere with the binding of IL-13 to IL-4R alpha-containing IL-13 receptor complexes. Antibodies may be screened, for example, in binding assays or tested for the ability to
20 inhibit an IL- IL-13-induced biological activity. An example of a suitable assay is illustrated in Example 2 below.

Antibodies specific for IL-4R alpha may be prepared using any technique known in the art. See, for example, *Monoclonal Antibodies, Hybridomas: A New Dimension in Biological Analyses*, Kennet *et al.* (eds.), Plenum Press, New York (1980); and *Antibodies: A Laboratory Manual*,
25 Harlow and Land (eds.), Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, (1988).

Antigen-binding fragments of antibodies of the invention may be produced by conventional techniques. Examples of such fragments include, but are not limited to, Fab and F(ab')₂ fragments. Antibody fragments and derivatives produced by genetic engineering techniques are also contemplated. Unless otherwise specified, the terms "antibody" and "monoclonal antibody"
30 as used herein encompass both whole antibodies and antigen-binding fragments and/or derivatives thereof.

Additional embodiments include chimeric antibodies, *e.g.*, humanized versions of murine monoclonal antibodies. Such humanized antibodies may be prepared by known techniques, and offer the advantage of reduced immunogenicity when the antibodies are administered to humans.
35 In one embodiment, a humanized monoclonal antibody comprises the variable domain of a murine antibody (or all or part of the antigen binding site thereof) and a constant domain derived from a human antibody. Alternatively, a humanized antibody fragment may comprise the antigen binding site of a murine monoclonal antibody and a variable domain fragment (lacking the antigen-binding site) derived from a human antibody. Procedures for the production of chimeric and further
40 engineered monoclonal antibodies include those described in Riechmann *et al.*, 1988, Nature 332:323, Liu *et al.*, 1987, Proc. Nat. Acad. Sci. USA 84:3439, Larrick *et al.*, 1989, Bio/Technology 7:934, and Winter *et al.*, 1993, TIPS 14:139. In one embodiment, the chimeric antibody is a CDR grafted antibody.

Procedures have been developed for generating human or partially human antibodies in
45 non-human animals. For example, mice in which one or more endogenous immunoglobulin genes

5 have been inactivated by various means have been prepared. Human immunoglobulin genes have been introduced into the mice to replace the inactivated mouse genes. Antibodies produced in the animal incorporate human immunoglobulin polypeptide chains encoded by the human genetic material introduced into the animal. In one embodiment, a non-human animal, such as a transgenic mouse, is immunized with an IL-4R polypeptide, such that antibodies directed against
10 the IL-4R polypeptide are generated in said animal. One example of a suitable immunogen is a soluble human IL-4R, such as a polypeptide comprising the extracellular domain of the protein of SEQ ID NO:2, or other immunogenic fragment of the protein of SEQ ID NO:2.

Examples of techniques for production and use of transgenic animals for the production of human or partially human antibodies are described in U.S. Patents 5,814,318, 5,569,825, and
15 5,545,806, which are incorporated by reference herein.

In another aspect, the present invention provides monoclonal antibodies that bind to IL-4 receptor. Monoclonal antibodies may be produced using any technique known in the art, *e.g.*, by immortalizing spleen cells harvested from the transgenic animal after completion of the immunization schedule. The spleen cells can be immortalized using any technique known in the
20 art, *e.g.*, by fusing them with myeloma cells to produce hybridomas. Myeloma cells for use in hybridoma-producing fusion procedures preferably are non-antibody-producing, have high fusion efficiency, and enzyme deficiencies that render them incapable of growing in certain selective media which support the growth of only the desired fused cells (hybridomas). Examples of suitable cell lines for use in mouse fusions include Sp-20, P3-X63/Ag8, P3-X63-Ag8.653,
25 NS1/1.Ag 4 1, Sp210-Ag14, FO, NSO/U, MPC-11, MPC11-X45-GTG 1.7 and S194/5XX0 Bul; examples of cell lines used in rat fusions include R210.RCY3, Y3-Ag 1.2.3, IR983F and 4B210. Other cell lines useful for cell fusions are U-266, GM1500-GRG2, LICR-LON-HMy2 and UC729-6.

In one embodiment, a hybridoma cell line is produced by immunizing an animal (*e.g.*, a transgenic animal having human immunoglobulin sequences) with an IL-4R immunogen;
30 harvesting spleen cells from the immunized animal; fusing the harvested spleen cells to a myeloma cell line, thereby generating hybridoma cells; establishing hybridoma cell lines from the hybridoma cells, and identifying a hybridoma cell line that produces an antibody that binds an IL-4R polypeptide. Such hybridoma cell lines, and anti-IL-4R monoclonal antibodies produced therefrom, are encompassed by the present invention.

35 Monoclonal antibodies secreted by a hybridoma cell line can be purified using any technique known in the art. Hybridomas or mAbs may be further screened to identify mAbs with particular properties, such as the ability to block an IL-4- and/or an IL-13-induced activity. Example of such screens are provided in Examples 2, 3, and 4.

Further examples of procedures for preparing antibodies directed against human IL-4
40 (including monoclonal antibodies), assays by which blocking antibodies are identified, and techniques for generating humanized or genetically engineered derivatives of anti-IL-4 antibodies, are described in U.S. Patents 5,041,381, 5,863,537, 5,928,904, and 5,676,940, which are hereby incorporated by reference. Further examples of antibodies that may be employed as IL-4 antagonists are described in WO 91/09059, also incorporated by reference herein.

5 In another aspect, the present invention provides human antibodies that bind IL-4R. In one embodiment of the invention, human antibodies raised against IL-4R alpha and produced by techniques involving use of transgenic mice, block binding of IL-4 and also IL-13 to cells. Such antibodies are IL-4 antagonists and additionally function as IL-13 antagonists.

10 Antibodies of the invention can comprise any constant region known in the art. The light chain constant region can be, for example, a kappa- or lambda-type light chain constant region, e.g., a human kappa- or lambda-type light chain constant region. The heavy chain constant region can be, for example, an alpha-, delta-, epsilon-, gamma-, or mu-type heavy chain constant regions, e.g., a human alpha-, delta-, epsilon-, gamma-, or mu-type heavy chain constant region. In one embodiment, the light or heavy chain constant region is a fragment, derivative, variant, or
15 mutein of a naturally occurring constant region.

Antibodies directed against an IL-4R can be used, for example, in assays to detect the presence of IL-4R polypeptides, either *in vitro* or *in vivo*. The antibodies also may be employed in purifying IL-4R proteins by immunoaffinity chromatography. Those antibodies that additionally can block binding of IL-4 to IL-4R may be used to inhibit a biological activity that results from such
20 binding. Blocking antibodies find use in the methods of the present invention. Such antibodies which function as IL-4 antagonists may be employed in treating any IL-4-induced condition, including but not limited to asthma and allergies, e.g., allergic rhinitis, contact dermatitis, and atopic dermatitis. In one embodiment, a human anti-IL-4R monoclonal antibody generated by procedures involving immunization of transgenic mice is employed in treating such conditions.

25 Antibodies may be employed in an *in vitro* procedure, or administered *in vivo* to inhibit an IL-4-induced biological activity. Disorders caused or exacerbated (directly or indirectly) by the interaction of IL-4 with cell surface IL-4 receptors, examples of which are provided above, thus may be treated. In one embodiment, the present invention provides a therapeutic method comprising *in vivo* administration of an IL-4 and/or IL-13 blocking antibody to a mammal in need
30 thereof in an amount effective for reducing an IL-4- and/or IL-13-induced biological activity.

Antibodies of the invention include, but are not limited to, partially human and fully human monoclonal antibodies that inhibit a biological activity of IL-4 and also inhibit a biological activity of IL-13. One embodiment is directed to a human monoclonal antibody that at least partially blocks binding of IL-4 to a cell, and at least partially blocks binding of IL-13 to a cell. In one embodiment,
35 the antibodies are generated by immunizing a transgenic mouse with an IL-4 receptor immunogen. In another embodiment, the immunogen is a human IL-4 receptor polypeptide. Hybridoma cell lines derived from the thus-immunized mice, wherein the hybridoma secretes a monoclonal antibody that binds IL-4R, also are provided herein.

Although human, partially human, or humanized antibodies will be suitable for many
40 applications, particularly those involving administration of the antibody to a human subject, other types of antibodies will be suitable for other applications. The non-human antibodies of the invention can be, for example, derived from any antibody-producing animal, such as mouse, rat, rabbit, goat, donkey, or non-human primate (such as monkey (e.g., cynomolgous or rhesus monkey) or ape (e.g., chimpanzee)). Non-human antibodies of the invention can be used, for
45 example, in *in vitro* and cell-culture based applications, or any other application where an immune

5 response to the antibody of the invention does not occur or is insignificant, can be prevented, is not a concern, or is desired. In one embodiment, a non-human antibody of the invention is administered to a non-human subject. In another embodiment, the non-human antibody does not elicit an immune response in the non-human subject. In another embodiment, the non-human antibody is from the same species as the non-human subject, *e.g.*, a mouse antibody of the invention is administered to a mouse. An antibody from a particular species can be made by, for example, immunizing an animal of that species with the desired immunogen (*e.g.*, a soluble IL-4 receptor polypeptide) or using an artificial system for generating antibodies of that species (*e.g.*, a bacterial or phage display-based system for generating antibodies of a particular species), or by converting an antibody from one species into an antibody from another species by replacing, *e.g.*, the constant region of the antibody with a constant region from the other species, or by replacing one or more amino acid residues of the antibody so that it more closely resembles the sequence of an antibody from the other species. In one embodiment, the antibody is a chimeric antibody comprising amino acid sequences derived from antibodies from two or more different species.

In another aspect, the present invention provides antibodies that comprise a light chain variable region selected from the group consisting of L1-L6 and/or a heavy chain variable region selected from the group consisting of H1-H24, and fragments, derivatives, muteins, and variants thereof (see Figures 2 and 3). Such an antibody can be denoted using the nomenclature "LxHy", wherein x corresponds to the number of the light chain variable region and y corresponds to the number of the heavy chain variable region as they are labeled in Figure 3. For example, L4H17 refers to an antibody with a light chain variable region comprising the amino acid sequence of L4 and a heavy chain variable region comprising the amino acid sequence of H17, as shown in Figure 3. Figures 2 and 3 also indicate the location of the CDR and framework regions of each of these variable domain sequences. Antibodies of the invention include, for example, L2H1, L3H1, L4H1, L5H1, L1H2, L1H3, L1H4, L1H5, L1H6, L1H7, L1H8, L1H9, L1H10, L1H11, L2H4, L2H12, L2H13, L2H14, L6H1, L2H2, L2H3, L2H6, L2H7, L2H8, L2H9, L2H10, and L2H11. Additional antibody variable sequences, *e.g.*, human antibody variable sequences, also can be used. See, *e.g.*, Sblattero *et al.*, 1998, *Immunotechnology* 3:271-78, de Haard *et al.*, 1999, *J. Biol. Chem.* 274:18218-30.

In one embodiment, the present invention provides an antibody comprising a light chain-variable domain comprising a sequence of amino acids that differs from the sequence of L1 only at one or more residues where any one of the sequences of L2-L6 differs from the sequence of L1 (*e.g.*, said sequence of said antibody differs from the sequence of L1 at residue(s) 1, 4, 7 *etc.*). In another embodiment, said sequence of said light chain-variable domain comprises at least one amino acid residue of any of the sequence of any one of L2-L6 at a position where it differs from the sequence of L1 (*e.g.*, said sequence comprises the residue(s) E1D, L4M, S7T, *etc.*). In another embodiment, said sequence differs from the sequence of L1 in at least one CDR (*e.g.*, CDR1, CDR2, or CDR3). In another embodiment, said sequence differs from the sequence of L1 in at least one FR (*e.g.*, FR1, FR2, FR3, or FR4). In another embodiment, the present invention provides an antibody comprising a heavy chain-variable domain comprising a sequence of amino acids that differs from the sequence of H1 only at one or more residues where any one of the

5 sequences of H2-H24 differs from the sequence of H1 (*e.g.*, said sequence of said antibody differs from the sequence of H1 at residue(s) 6, 13, 24 *etc.*). In another embodiment, said sequence of said heavy chain-variable domain comprises at least one amino acid residue of any of the sequence of any one of H2-H24 at a position where it differs from the sequence of H1 (*e.g.*, said sequence comprises the residue(s) Q6E, H13Q, G24A, *etc.*). In another embodiment, said
 10 sequence differs from the sequence of H1 in at least one CDR (*e.g.*, CDR2 or CDR3). In another embodiment, said sequence differs from the sequence of H1 in at least one FR (*e.g.*, FR1 or FR3).

In another embodiment, the present invention provides an antibody comprising an amino acid sequence selected from the group consisting of: a light chain complementarity determining
 15 region (CDR) 1 that differs by the insertion, deletion, or substitution of no more than 3, 2, 1, or 0 amino acid residues from a sequence selected from the group consisting of: residues 24-35 of SEQ ID NO:6, wherein the N at residue 8 is not substituted by an S; residues 24-35 of SEQ ID NO:8, wherein the T at residue 5 is not substituted by an S and/or the N at residue 7 is not substituted by an S and/or the D at residue 9 is not substituted by an S; residues 24-35 of SEQ ID
 20 NO:10, wherein the D at residue 9 is not substituted by an S; residues 24-35 of SEQ ID NO:12, wherein the N at residue 7 is not substituted by S and/or the N at residue 9 is not substituted by an S; and residues 24-35 of SEQ ID NO:14, wherein the G at residue 7 is not substituted by an S; a light chain CDR2 that differs by the insertion, deletion, or substitution of no more than 3, 2, 1, or 0 amino acid residues from a sequence selected from the group consisting of: residues 51-57 of
 25 SEQ ID NO:6, wherein the P at residue 7 is not substituted by a T; residues 51-57 of SEQ ID NO:10, wherein the S at residue 7 is not substituted by a T; and residues 51-57 of SEQ ID NO:12, wherein the T at residue 2 is not substituted by an A and/or the Y at residue 4 is not substituted by an S; a light chain CDR3 that differs by the insertion, deletion, or substitution of no more than 3, 2, 1, or 0 amino acid residues from a sequence selected from the group consisting of: residues 90-
 30 99 of SEQ ID NO:6, wherein the D at residue 4 is not substituted by a G, the H at residue 5 is not substituted by an S, the A at residue 7 is not substituted by a P, and/or the G at residue 8 is not substituted by a P; residues 90-99 of SEQ ID NO:8, wherein the R at residue 5 is not substituted by an S; and residues 90-99 of SEQ ID NO:14, wherein the M a residue 10 is not substituted by a T; a heavy chain CDR2 that differs by the insertion, deletion, or substitution of no more than 3
 35 amino acid residues from the sequence of residues 50-65 of SEQ ID NO:18, wherein the S at residue 9 is not substituted by an N; and a heavy chain CDR3 that differs by the insertion, deletion, or substitution of no more than 3, 2, 1, or 0 amino acid residues from a sequence selected from the group consisting of: residues 98-104 of SEQ ID NO:18, wherein the T at residue 6 is not substituted by a D and/or the H at residue 7 is not substituted by a Y; residues 98-104 of SEQ ID
 40 NO:20, wherein the W at residue 4 is not substituted by a Y, the Y at residue 5 is not substituted by an F, the N at residue 6 is not substituted by a D, and/or the N at residue 7 is not substituted by a Y; residues 98-104 of SEQ ID NO:22, wherein the P at residue 6 is not substituted by a D and/or the W at residue 7 is not substituted by a Y; residues 98-104 of SEQ ID NO:24, wherein the T at residue 6 is not substituted by a D and/or the R at position 7 is not substituted by a Y; residues 98-

5 104 of SEQ ID NO:26, wherein the Y at residue 5 is not substituted by an F; residues 98-104 of SEQ ID NO:30, wherein the W at residue 4 is not substituted by a Y; and residues 98-104 of SEQ ID NO:34, wherein the W at residue 4 is not substituted by a Y and/or the Y at residue 5 is not substituted by an F; wherein said antibody binds to IL-4 receptor alpha.

In another embodiment, the antibody inhibits the binding of IL-4 to an IL-4 receptor. In another embodiment, the antibody inhibits the binding of IL-13 to an IL-13 receptor. In another embodiment, the antibody inhibits the binding of IL-4 to an IL-4 receptor and of IL-13 to an IL-13 receptor. In another embodiment, the antibody specifically binds IL-4 receptor alpha.

In another embodiment, the antibody comprises two or three such light chain complementarity determining regions (CDR).

15 In another embodiment, the antibody comprises two or three such heavy chain complementarity determining regions.

In another embodiment, the antibody further comprises a framework segment (FR) comprising a sequence selected from the group consisting of: a light chain framework segment (FR) 1 that differs by the insertion, deletion, or substitution of no more than 3, 2, 1, or 0 amino acid residues from a sequence selected from the group consisting of: residues 1-23 of SEQ ID NO:4; residues 1-23 of SEQ ID NO:10; residues 1-23 of SEQ ID NO:12; and residues 1-23 of SEQ ID NO:14; a light chain FR2 that differs by the insertion, deletion, or substitution of no more than 3, 2, 1, or 0 amino acid residues from a sequence selected from the group consisting of: residues 36-50 of SEQ ID NO:4; residues 36-50 of SEQ ID NO:6; and residues 36-50 of SEQ ID NO:14; a light chain FR3 that differs by the insertion, deletion, or substitution of no more than 3, 2, 1, or 0 amino acid residues from a sequence selected from the group consisting of: residues 58-89 of SEQ ID NO:4; residues 58-89 of SEQ ID NO:10; and residues 58-89 of SEQ ID NO:12; a light chain FR4 that differs by the insertion, deletion, or substitution of no more than 3, 2, 1, or 0 amino acid residues from a sequence selected from the group consisting of: residues 100-109 of SEQ ID NO:4; residues 100-109 of SEQ ID NO:8; and residues 100-109 of SEQ ID NO:12; a heavy chain FR1 that differs by the insertion, deletion, or substitution of no more than 3, 2, 1, or 0 amino acid residues from a sequence selected from the group consisting of: residues 1-30 of SEQ ID NO:16; and residues 1-30 of SEQ ID NO:42; heavy chain FR2 that differs by the insertion, deletion, or substitution of no more than 3, 2, 1, or 0 amino acid residues from the sequence residues 36-49 of SEQ ID NO:16, a heavy chain FR3 that differs by the insertion, deletion, or substitution of no more than 3, 2, 1, or 0 amino acid residues from a sequence selected from the group consisting of: residues 66-97 of SEQ ID NO:16; residues 66-97 of SEQ ID NO:18; and residues 66-97 of SEQ ID NO:42; and a heavy chain FR4 that differs by the insertion, deletion, or substitution of no more than 3 amino acid residues from the sequence residues 105-115 of SEQ ID NO:16.

40 In another embodiment, the antibody comprises a light chain variable domain that is at least 80, 85, 90, 95, or 100% identical to a sequence selected from the group consisting of: SEQ ID NO:6, 8, 10, 12, and 14, with the proviso that the said light chain variable domain does not comprise the sequence of SEQ ID NO:4.

5 In another embodiment, the antibody comprises a heavy chain variable domain that is at least 80, 85, 90, 95, or 100% identical to a sequence selected from the group consisting of: SEQ ID NO:18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 46, 48, 50, 52, 54, 56, 58, 60, and 62 with the proviso that said heavy chain variable domain does not comprise the sequence of SEQ ID NO:16.

10 In another embodiment, the present invention provides a fragment of such an antibody, wherein said fragment binds to IL-4 receptor alpha.

In one embodiment, particular antibodies of the invention are selected from the group consisting of L2H1; an antibody that is cross-reactive L2H1, an antibody that binds to the same epitope as L2H1; an antibody that competes with L2H1 for binding to a cell that expresses human
15 IL-4R; an antibody that possesses a biological activity of L2H1 and an antigen-binding fragment (including one derived by recombinant means) of L2H1. In one embodiment, the antibody has a binding affinity for human IL-4R that is substantially equivalent to the binding affinity of L2H1 for human IL-4R. Hybridoma cell lines that produce any such antibodies also are provided by the present invention.

20 One example of a biological activity of L2H1 is the ability to function as both an IL-4 antagonist and an IL-13 antagonist. In one embodiment, an antibody of the invention possesses IL-4-blocking activity substantially equivalent to that of L2H1; and possesses IL-13-blocking activity substantially equivalent to that of L2H1. Such activity may be measured in any suitable conventional assay (e.g., as measured in the CD23 expression assay described in Example 2).

25 Complementarity determining regions (CDRs) of a given antibody may be identified using the system described by Kabat *et al.* in *Sequences of Proteins of Immunological Interest*, 5th Ed., US Dept. of Health and Human Services, PHS, NIH, NIH Publication no. 91-3242, 1991. Particular embodiments of antibodies of the present invention comprise, within the variable domain of their light chain, at least one of the complementarity determining regions (CDRs), or hypervariable
30 regions, found in the light chain of L2H1. CDRs of L2H1 are discussed in Example 5. Thus, among the antibodies provided herein are those comprising from one to all three of the following sequences in the light chain variable domain of L2H1: amino acid residues 24-35; residues 51-57; and residues 90-99 of SEQ ID NO:6. Particular antibodies provided herein comprise, within the variable domain of their heavy chain, at least one of the CDRs found in the heavy chain of L2H1.
35 Thus, among the antibodies provided herein are those comprising from one to all three of the following sequences in the heavy chain variable domain: residues 31-35; residues 50-65; and residues 98-104 of SEQ ID NO:16.

The DNA sequence of the variable domain of the light chain of L2H1 is presented in SEQ ID NO:5, and the encoded amino acid sequence is presented in SEQ ID NO:6. The DNA
40 sequence for the variable domain of the heavy chain of L2H1 is presented as SEQ ID NO:15, and the encoded amino acid sequence is presented in SEQ ID NO:16. Antibodies of the present invention include, but are not limited to, antibodies that comprise, in their light chain, SEQ ID NO:6; and antibodies that additionally or alternatively comprise, in their heavy chain, SEQ ID NO:16.

5 In another embodiment, particular antibodies of the invention are selected from the group consisting of L3H1; an antibody that is cross-reactive with L3H1; an antibody that binds to the same epitope as L3H1; an antibody that competes with L3H1 for binding to a cell that expresses human IL-4R; an antibody that possesses a biological activity of L3H1; and an antigen-binding
10 fragment of any of the foregoing antibodies. In one embodiment, the antibody has a binding affinity for human IL-4R that is substantially equivalent to the binding affinity of L3H1 for human IL-4R. Hybridoma cell lines that produce any such antibodies also are provided by the present invention.

One example of a biological activity of L3H1 is the ability to function as both an IL-4 antagonist and an IL-13 antagonist. In one embodiment, an antibody of the invention possesses
15 IL-4-blocking activity substantially equivalent to that of L3H1, and possesses IL-13-blocking activity substantially equivalent to that of L3H1. Such activity may be measured in any suitable conventional assay (*e.g.*, as measured in the CD23 expression assay described in Example 2).

IgG4 antibodies derived from L3H1 are provided herein. Another embodiment is directed to IgM antibodies derived from L3H1. Procedures for switching (altering) the subclass or isotype
20 of an antibody are known in the pertinent field. Such procedures may involve, for example, recombinant DNA technology, whereby DNA encoding antibody polypeptide chains that confer the desired subclass is substituted for DNA encoding the corresponding polypeptide chain of the parent antibody.

The DNA sequence of the variable domain of the light chain of L3H1 is presented in SEQ
25 ID NO:7, and the encoded amino acid sequence is presented in SEQ ID NO:8. The DNA sequence for the variable domain of the heavy chain of L3H1 is presented as SEQ ID NO:15, and the encoded amino acid sequence is presented in SEQ ID NO:16. Antibodies of the present invention include, but are not limited to, antibodies that comprise, in their light chain, residues 1 to 109 of SEQ ID NO:8; and antibodies that additionally or alternatively comprise, in their heavy
30 chain, residues 1 to 115 of SEQ ID NO:16.

Particular embodiments of antibodies of the present invention comprise, within the variable domain of their light chain, at least one of the complementarity determining regions (CDRs), or hypervariable regions, found in the light chain of L3H1. CDRs of L3H1 are discussed
35 in example 5. Thus, among the antibodies provided herein are those comprising from one to all three of the following sequences in the light chain variable domain: amino acid residues 24-35; residues 51-57; and residues 90-99 of SEQ ID NO:8. Particular antibodies provided herein comprise, within the variable domain of their heavy chain, at least one of the CDRs found in the heavy chain of L3H1. Thus, among the antibodies provided herein are those comprising from one to all three of the following sequences in the heavy chain variable domain: residues 31-35;
40 residues 50-65; and residues 98-104 of SEQ ID NO:16.

In another embodiment, particular antibodies of the invention are selected from the group consisting of L4H1; an antibody that is cross-reactive with L4H1; an antibody that binds to the same epitope as L4H1; an antibody that competes with L4H1 for binding to a cell that expresses human IL-4R; an antibody that possesses a biological activity of L4H1; and an antigen-binding
45 fragment of any of the foregoing antibodies. In one embodiment, the antibody has a binding

5 affinity for human IL-4R that is substantially equivalent to the binding affinity of L4H1 for human IL-4R. Hybridoma cell lines that produce any such antibodies also are provided by the present invention.

One example of a biological activity of L4H1 is the ability to function as both an IL-4 antagonist and an IL-13 antagonist. In one embodiment, an antibody of the invention possesses
10 IL-4-blocking activity substantially equivalent to that of L4H1; and possesses IL-13-blocking activity substantially equivalent to that of L4H1. Such activity may be measured in any suitable conventional assay (*e.g.*, as measured in the CD23 expression assay described in Example 2).

The DNA sequence of the variable domain of the light chain of L4H1 is presented in SEQ ID NO:9, and the encoded amino acid sequence is presented in SEQ ID NO:10. The DNA
15 sequence for the variable domain of the heavy chain of L4H1 is presented as SEQ ID NO:15, and the encoded amino acid sequence is presented in SEQ ID NO:16. Antibodies of the present invention include, but are not limited to, antibodies that comprise, in their light chain, residues 1 to 109 of SEQ ID NO:10; and antibodies that additionally or alternatively comprise, in their heavy chain, residues 1 to 115 of SEQ ID NO:16.

20 Particular embodiments of antibodies of the present invention comprise, within the variable domain of their light chain, at least one of the complementarity determining regions (CDRs), or hypervariable regions, found in the light chain of L4H1. CDRs of L4H1 are discussed in example 5. Thus, among the antibodies provided herein are those comprising from one to all three of the following sequences in the light chain variable domain: amino acid residues 24-35;
25 residues 51-57; and residues 90-99 of SEQ ID NO:10. Particular antibodies provided herein comprise, within the variable domain of their heavy chain, at least one of the CDRs found in the heavy chain of L4H1. Thus, among the antibodies provided herein are those comprising from one to all three of the following sequences in the heavy chain variable domain: residues 31-35; residues 50-65; and residues 99-104 of SEQ ID NO:16.

30 In another embodiment, particular antibodies of the invention are selected from the group consisting of L5H1; an antibody that is cross-reactive with L5H1; an antibody that binds to the same epitope as L5H1; an antibody that competes with L5H1 for binding to a cell that expresses human IL-4R; an antibody that possesses a biological activity of L5H1; and an antigen-binding fragment of any of the foregoing antibodies. In one embodiment, the antibody has a binding
35 affinity for human IL-4R that is substantially equivalent to the binding affinity of L5H1 for human IL-4R. Hybridoma cell lines that produce any such antibodies also are provided by the present invention.

One example of a biological activity of L5H1 is the ability to function as both an IL-4 antagonist and an IL-13 antagonist. In one embodiment, an antibody of the invention possesses
40 IL-4-blocking activity substantially equivalent to that of L5H1; and possesses IL-13-blocking activity substantially equivalent to that of L5H1. Such activity may be measured in any suitable conventional assay (*e.g.*, as measured in the CD23 expression assay described in Example 2).

The DNA sequence of the variable domain of the light chain of L5H1 is presented in SEQ ID NO:11, and the encoded amino acid sequence is presented in SEQ ID NO:12. The DNA
45 sequence for the variable domain of the heavy chain of L5H1 is presented as SEQ ID NO:15, and

5 the encoded amino acid sequence is presented in SEQ ID NO:16. Antibodies of the present invention include, but are not limited to, antibodies that comprise, in their light chain, residues 1 to 109 of SEQ ID NO:12; and antibodies that additionally or alternatively comprise, in their heavy chain, residues 1 to 115 of SEQ ID NO:16.

10 Particular embodiments of antibodies of the present invention comprise, within the variable domain of their light chain, at least one of the complementarity determining regions (CDRs), or hypervariable regions, found in the light chain of L5H1. CDRs of L5H1 are discussed in example 5. Thus, among the antibodies provided herein are those comprising from one to all three of the following sequences in the light chain variable domain: amino acid residues 24-35; residues 51-57; and residues 90-99 of SEQ ID NO:12. Particular antibodies provided herein
15 comprise, within the variable domain of their heavy chain, at least one of the CDRs found in the heavy chain of L5H1. Thus, among the antibodies provided herein are those comprising from one to all three of the following sequences in the heavy chain variable domain: residues 31-35; residues 50-65; and residues 98-104 of SEQ ID NO:16.

20 In another embodiment, particular antibodies of the invention are selected from the group consisting of L1H2; an antibody that is cross-reactive with L1H2; an antibody that binds to the same epitope as L1H2; an antibody that competes with L1H2 for binding to a cell that expresses human IL-4R; an antibody that possesses a biological activity of L1H2; and an antigen-binding fragment of any of the foregoing antibodies. In one embodiment, the antibody has a binding affinity for human IL-4R that is substantially equivalent to the binding affinity of L1H2 for human IL-
25 4R. Hybridoma cell lines that produce any such antibodies also are provided by the present invention.

One example of a biological activity of L1H2 is the ability to function as both an IL-4 antagonist and an IL-13 antagonist. In one embodiment, an antibody of the invention possesses IL-4-blocking activity substantially equivalent to that of L1H2; and possesses IL-13-blocking activity
30 substantially equivalent to that of L1H2. Such activity may be measured in any suitable conventional assay (e.g., as measured in the CD23 expression assay described in Example 2).

The DNA sequence of the variable domain of the light chain of L1H2 is presented in SEQ ID NO:3, and the encoded amino acid sequence is presented in SEQ ID NO:4. The DNA sequence for the variable domain of the heavy chain of L1H2 is presented as SEQ ID NO:17, and
35 the encoded amino acid sequence is presented in SEQ ID NO:18. Antibodies of the present invention include, but are not limited to, antibodies that comprise, in their light chain, residues 1 to 109 of SEQ ID NO:4; and antibodies that additionally or alternatively comprise, in their heavy chain, residues 1 to 115 of SEQ ID NO:18.

40 Particular embodiments of antibodies of the present invention comprise, within the variable domain of their light chain, at least one of the complementarity determining regions (CDRs), or hypervariable regions, found in the light chain of L1H2. CDRs of L1H2 are discussed in example 5. Thus, among the antibodies provided herein are those comprising from one to all three of the following sequences in the light chain variable domain: amino acid residues 24-35; residues 51-57; and residues 90-99 of SEQ ID NO:4. Particular antibodies provided herein
45 comprise, within the variable domain of their heavy chain, at least one of the CDRs found in the

5 heavy chain of L1H2. Thus, among the antibodies provided herein are those comprising from one to all three of the following sequences in the heavy chain variable domain: residues 31-35; residues 50-65; and residues 98-104 of SEQ ID NO:18.

In another embodiment, particular antibodies of the invention are selected from the group consisting of L1H3; an antibody that is cross-reactive with L1H3; an antibody that binds to the
10 same epitope as L1H3; an antibody that competes with L1H3 for binding to a cell that expresses human IL-4R; an antibody that possesses a biological activity of L1H3; and an antigen-binding fragment of any of the foregoing antibodies. In one embodiment, the antibody has a binding affinity for human IL-4R that is substantially equivalent to the binding affinity of L1H3 for human IL-4R. Hybridoma cell lines that produce any such antibodies also are provided by the present
15 invention.

One example of a biological activity of L1H3 is the ability to function as both an IL-4 antagonist and an IL-13 antagonist. In one embodiment, an antibody of the invention possesses IL-4-blocking activity substantially equivalent to that of L1H3; and possesses IL-13-blocking activity substantially equivalent to that of L1H3. Such activity may be measured in any suitable
20 conventional assay (e.g., as measured in the CD23 expression assay described in Example 2).

The DNA sequence of the variable domain of the light chain of L1H3 is presented in SEQ ID NO:3, and the encoded amino acid sequence is presented in SEQ ID NO:4. The DNA sequence for the variable domain of the heavy chain of L1H3 is presented as SEQ ID NO:19, and the encoded amino acid sequence is presented in SEQ ID NO:20. Antibodies of the present
25 invention include, but are not limited to, antibodies that comprise, in their light chain, residues 1 to 109 of SEQ ID NO:4; and antibodies that additionally or alternatively comprise, in their heavy chain, residues 1 to 115 of SEQ ID NO:20.

Particular embodiments of antibodies of the present invention comprise, within the variable domain of their light chain, at least one of the complementarity determining regions
30 (CDRs), or hypervariable regions, found in the light chain of L1H3. CDRs of L1H3 are discussed in example 5. Thus, among the antibodies provided herein are those comprising from one to all three of the following sequences in the light chain variable domain: amino acid residues 24-35; residues 51-57; and residues 90-99 of SEQ ID NO:4. Particular antibodies provided herein comprise, within the variable domain of their heavy chain, at least one of the CDRs found in the
35 heavy chain of L1H3. Thus, among the antibodies provided herein are those comprising from one to all three of the following sequences in the heavy chain variable domain: residues 31-35; residues 50-65; and residues 98-104 of SEQ ID NO:20.

In another embodiment, particular antibodies of the invention are selected from the group consisting of L1H4; an antibody that is cross-reactive with L1H4; an antibody that binds to the
40 same epitope as L1H4; an antibody that competes with L1H4 for binding to a cell that expresses human IL-4R; an antibody that possesses a biological activity of L1H4; and an antigen-binding fragment of any of the foregoing antibodies. In one embodiment, the antibody has a binding affinity for human IL-4R that is substantially equivalent to the binding affinity of L1H4 for human IL-4R. Hybridoma cell lines that produce any such antibodies also are provided by the present
45 invention.

5 One example of a biological activity of L1H4 is the ability to function as both an IL-4 antagonist and an IL-13 antagonist. In one embodiment, an antibody of the invention possesses IL-4-blocking activity substantially equivalent to that of L1H4; and possesses IL-13-blocking activity substantially equivalent to that of L1H4. Such activity may be measured in any suitable conventional assay (*e.g.*, as measured in the CD23 expression assay described in Example 2).

10 The DNA sequence of the variable domain of the light chain of L1H4 is presented in SEQ ID NO:3, and the encoded amino acid sequence is presented in SEQ ID NO:4. The DNA sequence for the variable domain of the heavy chain of L1H4 is presented as SEQ ID NO:21, and the encoded amino acid sequence is presented in SEQ ID NO:22. Antibodies of the present invention include, but are not limited to, antibodies that comprise, in their light chain, residues 1 to
15 109 of SEQ ID NO:4; and antibodies that additionally or alternatively comprise, in their heavy chain, residues 1 to 115 of SEQ ID NO:22.

Particular embodiments of antibodies of the present invention comprise, within the variable domain of their light chain, at least one of the complementarity determining regions (CDRs), or hypervariable regions, found in the light chain of L1H4. CDRs of L1H4 are discussed
20 in example 5. Thus, among the antibodies provided herein are those comprising from one to all three of the following sequences in the light chain variable domain: amino acid residues 24-35; residues 51-57; and residues 90-99 of SEQ ID NO:4. Particular antibodies provided herein comprise, within the variable domain of their heavy chain, at least one of the CDRs found in the heavy chain of L1H4. Thus, among the antibodies provided herein are those comprising from one
25 to all three of the following sequences in the heavy chain variable domain: residues 31-35; residues 50-65; and residues 98-104 of SEQ ID NO:22.

In another embodiment, particular antibodies of the invention are selected from the group consisting of L1H5; an antibody that is cross-reactive with L1H5; an antibody that binds to the same epitope as L1H5; an antibody that competes with L1H5 for binding to a cell that expresses
30 human IL-4R; an antibody that possesses a biological activity of L1H5; and an antigen-binding fragment of any of the foregoing antibodies. In one embodiment, the antibody has a binding affinity for human IL-4R that is substantially equivalent to the binding affinity of L1H5 for human IL-4R. Hybridoma cell lines that produce any such antibodies also are provided by the present invention.

35 One example of a biological activity of L1H5 is the ability to function as both an IL-4 antagonist and an IL-13 antagonist. In one embodiment, an antibody of the invention possesses IL-4-blocking activity substantially equivalent to that of L1H5; and possesses IL-13-blocking activity substantially equivalent to that of L1H5. Such activity may be measured in any suitable conventional assay (*e.g.*, as measured in the CD23 expression assay described in Example 2).

40 The DNA sequence of the variable domain of the light chain of L1H5 is presented in SEQ ID NO:3, and the encoded amino acid sequence is presented in SEQ ID NO:4. The DNA sequence for the variable domain of the heavy chain of L1H5 is presented as SEQ ID NO:23, and the encoded amino acid sequence is presented in SEQ ID NO:24. Antibodies of the present invention include, but are not limited to, antibodies that comprise, in their light chain, residues 1 to

5 109 of SEQ ID NO:4; and antibodies that additionally or alternatively comprise, in their heavy chain, residues 1 to 115 of SEQ ID NO:24.

Particular embodiments of antibodies of the present invention comprise, within the variable domain of their light chain, at least one of the complementarity determining regions (CDRs), or hypervariable regions, found in the light chain of L1H5. CDRs of L1H5 are discussed
10 in example 5. Thus, among the antibodies provided herein are those comprising from one to all three of the following sequences in the light chain variable domain: amino acid residues 24-35; residues 51-57; and residues 90-99 of SEQ ID NO:4. Particular antibodies provided herein comprise, within the variable domain of their heavy chain, at least one of the CDRs found in the heavy chain of L1H5. Thus, among the antibodies provided herein are those comprising from one
15 to all three of the following sequences in the heavy chain variable domain: residues 31-35; residues 50-65; and residues 98-104 of SEQ ID NO:24.

In another embodiment, particular antibodies of the invention are selected from the group consisting of L1H6; an antibody that is cross-reactive with L1H6; an antibody that binds to the same epitope as L1H6; an antibody that competes with L1H6 for binding to a cell that expresses
20 human IL-4R; an antibody that possesses a biological activity of L1H6; and an antigen-binding fragment of any of the foregoing antibodies. In one embodiment, the antibody has a binding affinity for human IL-4R that is substantially equivalent to the binding affinity of L1H6 for human IL-4R. Hybridoma cell lines that produce any such antibodies also are provided by the present invention.

25 One example of a biological activity of L1H6 is the ability to function as both an IL-4 antagonist and an IL-13 antagonist. In one embodiment, an antibody of the invention possesses IL-4-blocking activity substantially equivalent to that of L1H6; and possesses IL-13-blocking activity substantially equivalent to that of L1H6. Such activity may be measured in any suitable conventional assay (e.g., as measured in the CD23 expression assay described in Example 2).

30 The DNA sequence of the variable domain of the light chain of L1H6 is presented in SEQ ID NO:3, and the encoded amino acid sequence is presented in SEQ ID NO:4. The DNA sequence for the variable domain of the heavy chain of L1H6 is presented as SEQ ID NO:25, and the encoded amino acid sequence is presented in SEQ ID NO:26. Antibodies of the present invention include, but are not limited to, antibodies that comprise, in their light chain, residues 1 to
35 109 of SEQ ID NO:4; and antibodies that additionally or alternatively comprise, in their heavy chain, residues 1 to 115 of SEQ ID NO:26.

Particular embodiments of antibodies of the present invention comprise, within the variable domain of their light chain, at least one of the complementarity determining regions (CDRs), or hypervariable regions, found in the light chain of L1H6. CDRs of L1H6 are discussed
40 in example 5. Thus, among the antibodies provided herein are those comprising from one to all three of the following sequences in the light chain variable domain: amino acid residues 24-35; residues 51-57; and residues 90-99 of SEQ ID NO:4. Particular antibodies provided herein comprise, within the variable domain of their heavy chain, at least one of the CDRs found in the heavy chain of L1H6. Thus, among the antibodies provided herein are those comprising from one

5 to all three of the following sequences in the heavy chain variable domain: residues 31-35; residues 50-65; and residues 98-104 of SEQ ID NO:26.

In another embodiment, particular antibodies of the invention are selected from the group consisting of L1H7; an antibody that is cross-reactive with L1H7; an antibody that binds to the same epitope as L1H7; an antibody that competes with L1H7 for binding to a cell that expresses
10 human IL-4R; an antibody that possesses a biological activity of L1H7; and an antigen-binding fragment of any of the foregoing antibodies. In one embodiment, the antibody has a binding affinity for human IL-4R that is substantially equivalent to the binding affinity of L1H7 for human IL-4R. Hybridoma cell lines that produce any such antibodies also are provided by the present invention.

15 One example of a biological activity of L1H7 is the ability to function as both an IL-4 antagonist and an IL-13 antagonist. In one embodiment, an antibody of the invention possesses IL-4-blocking activity substantially equivalent to that of L1H7; and possesses IL-13-blocking activity substantially equivalent to that of L1H7. Such activity may be measured in any suitable conventional assay (e.g., as measured in the CD23 expression assay described in Example 2).

20 The DNA sequence of the variable domain of the light chain of L1H7 is presented in SEQ ID NO:3, and the encoded amino acid sequence is presented in SEQ ID NO:4. The DNA sequence for the variable domain of the heavy chain of L1H7 is presented as SEQ ID NO:27, and the encoded amino acid sequence is presented in SEQ ID NO:28. Antibodies of the present invention include, but are not limited to, antibodies that comprise, in their light chain, residues 1 to
25 109 of SEQ ID NO:4; and antibodies that additionally or alternatively comprise, in their heavy chain, residues 1 to 115 of SEQ ID NO:28.

Particular embodiments of antibodies of the present invention comprise, within the variable domain of their light chain, at least one of the complementarity determining regions (CDRs), or hypervariable regions, found in the light chain of L1H7. CDRs of L1H7 are discussed
30 in example 5. Thus, among the antibodies provided herein are those comprising from one to all three of the following sequences in the light chain variable domain: amino acid residues 24-35; residues 51-57; and residues 90-99 of SEQ ID NO:4. Particular antibodies provided herein comprise, within the variable domain of their heavy chain, at least one of the CDRs found in the heavy chain of L1H7. Thus, among the antibodies provided herein are those comprising from one
35 to all three of the following sequences in the heavy chain variable domain: residues 31-35; residues 50-65; and residues 98-104 of SEQ ID NO:28.

In another embodiment, particular antibodies of the invention are selected from the group consisting of L1H8; an antibody that is cross-reactive with L1H8; an antibody that binds to the same epitope as L1H8; an antibody that competes with L1H8 for binding to a cell that expresses
40 human IL-4R; an antibody that possesses a biological activity of L1H8; and an antigen-binding fragment of any of the foregoing antibodies. In one embodiment, the antibody has a binding affinity for human IL-4R that is substantially equivalent to the binding affinity of L1H8 for human IL-4R. Hybridoma cell lines that produce any such antibodies also are provided by the present invention.

5 One example of a biological activity of L1H8 is the ability to function as both an IL-4 antagonist and an IL-13 antagonist. In one embodiment, an antibody of the invention possesses IL-4-blocking activity substantially equivalent to that of L1H8; and possesses IL-13-blocking activity substantially equivalent to that of L1H8. Such activity may be measured in any suitable conventional assay (e.g., as measured in the CD23 expression assay described in Example 2).

10 The DNA sequence of the variable domain of the light chain of L1H8 is presented in SEQ ID NO:3, and the encoded amino acid sequence is presented in SEQ ID NO:4. The DNA sequence for the variable domain of the heavy chain of L1H8 is presented as SEQ ID NO:29, and the encoded amino acid sequence is presented in SEQ ID NO:30. Antibodies of the present invention include, but are not limited to, antibodies that comprise, in their light chain, residues 1 to
15 109 of SEQ ID NO:4; and antibodies that additionally or alternatively comprise, in their heavy chain, residues 1 to 115 of SEQ ID NO:30.

 Particular embodiments of antibodies of the present invention comprise, within the variable domain of their light chain, at least one of the complementarity determining regions (CDRs), or hypervariable regions, found in the light chain of L1H8. CDRs of L1H8 are discussed
20 in example 5. Thus, among the antibodies provided herein are those comprising from one to all three of the following sequences in the light chain variable domain: amino acid residues 24-35; residues 51-57; and residues 90-99 of SEQ ID NO:4. Particular antibodies provided herein comprise, within the variable domain of their heavy chain, at least one of the CDRs found in the heavy chain of L1H8. Thus, among the antibodies provided herein are those comprising from one
25 to all three of the following sequences in the heavy chain variable domain: residues 31-35; residues 50-65; and residues 98-104 of SEQ ID NO:30.

 In another embodiment, particular antibodies of the invention are selected from the group consisting of L1H9; an antibody that is cross-reactive with L1H9; an antibody that binds to the same epitope as L1H9; an antibody that competes with L1H9 for binding to a cell that expresses
30 human IL-4R; an antibody that possesses a biological activity of L1H9; and an antigen-binding fragment of any of the foregoing antibodies. In one embodiment, the antibody has a binding affinity for human IL-4R that is substantially equivalent to the binding affinity of L1H9 for human IL-4R. Hybridoma cell lines that produce any such antibodies also are provided by the present invention.

35 One example of a biological activity of L1H9 is the ability to function as both an IL-4 antagonist and an IL-13 antagonist. In one embodiment, an antibody of the invention possesses IL-4-blocking activity substantially equivalent to that of L1H9; and possesses IL-13-blocking activity substantially equivalent to that of L1H9. Such activity may be measured in any suitable conventional assay (e.g., as measured in the CD23 expression assay described in Example 2).

40 The DNA sequence of the variable domain of the light chain of L1H9 is presented in SEQ ID NO:3, and the encoded amino acid sequence is presented in SEQ ID NO:4. The DNA sequence for the variable domain of the heavy chain of L1H9 is presented as SEQ ID NO:31, and the encoded amino acid sequence is presented in SEQ ID NO:32. Antibodies of the present invention include, but are not limited to, antibodies that comprise, in their light chain, residues 1 to

5 109 of SEQ ID NO:4; and antibodies that additionally or alternatively comprise, in their heavy chain, residues 1 to 115 of SEQ ID NO:32.

Particular embodiments of antibodies of the present invention comprise, within the variable domain of their light chain, at least one of the complementarity determining regions (CDRs), or hypervariable regions, found in the light chain of L1H9. CDRs of L1H9 are discussed
10 in example 5. Thus, among the antibodies provided herein are those comprising from one to all three of the following sequences in the light chain variable domain: amino acid residues 24-35; residues 51-57; and residues 90-99 of SEQ ID NO:4. Particular antibodies provided herein comprise, within the variable domain of their heavy chain, at least one of the CDRs found in the heavy chain of L1H9. Thus, among the antibodies provided herein are those comprising from one
15 to all three of the following sequences in the heavy chain variable domain: residues 31-35; residues 50-65; and residues 98-104 of SEQ ID NO:32.

In another embodiment, particular antibodies of the invention are selected from the group consisting of L1H10; an antibody that is cross-reactive with L1H10; an antibody that binds to the same epitope as L1H10; an antibody that competes with L1H10 for binding to a cell that expresses
20 human IL-4R; an antibody that possesses a biological activity of L1H10; and an antigen-binding fragment of any of the foregoing antibodies. In one embodiment, the antibody has a binding affinity for human IL-4R that is substantially equivalent to the binding affinity of L1H10 for human IL-4R. Hybridoma cell lines that produce any such antibodies also are provided by the present invention.

25 One example of a biological activity of L1H10 is the ability to function as both an IL-4 antagonist and an IL-13 antagonist. In one embodiment, an antibody of the invention possesses IL-4-blocking activity substantially equivalent to that of L1H10; and possesses IL-13-blocking activity substantially equivalent to that of L1H10. Such activity may be measured in any suitable conventional assay (e.g., as measured in the CD23 expression assay described in Example 2).

30 The DNA sequence of the variable domain of the light chain of L1H10 is presented in SEQ ID NO:3, and the encoded amino acid sequence is presented in SEQ ID NO:4. The DNA sequence for the variable domain of the heavy chain of L1H10 is presented as SEQ ID NO:33, and the encoded amino acid sequence is presented in SEQ ID NO:34. Antibodies of the present invention include, but are not limited to, antibodies that comprise, in their light chain, residues 1 to
35 109 of SEQ ID NO:4; and antibodies that additionally or alternatively comprise, in their heavy chain, residues 1 to 115 of SEQ ID NO:34.

Particular embodiments of antibodies of the present invention comprise, within the variable domain of their light chain, at least one of the complementarity determining regions (CDRs), or hypervariable regions, found in the light chain of L1H10. CDRs of L1H10 are
40 discussed in example 5. Thus, among the antibodies provided herein are those comprising from one to all three of the following sequences in the light chain variable domain: amino acid residues 24-35; residues 51-57; and residues 90-99 of SEQ ID NO:4. Particular antibodies provided herein comprise, within the variable domain of their heavy chain, at least one of the CDRs found in the heavy chain of L1H10. Thus, among the antibodies provided herein are those comprising from

- 5 one to all three of the following sequences in the heavy chain variable domain: residues 31-35; residues 50-65; and residues 98-104 of SEQ ID NO:34.

In another embodiment, particular antibodies of the invention are selected from the group consisting of L1H11; an antibody that is cross-reactive with L1H11; an antibody that binds to the same epitope as L1H11; an antibody that competes with L1H11 for binding to a cell that expresses
10 human IL-4R; an antibody that possesses a biological activity of L1H11; and an antigen-binding fragment of any of the foregoing antibodies. In one embodiment, the antibody has a binding affinity for human IL-4R that is substantially equivalent to the binding affinity of L1H11 for human IL-4R. Hybridoma cell lines that produce any such antibodies also are provided by the present invention.

15 One example of a biological activity of L1H11 is the ability to function as both an IL-4 antagonist and an IL-13 antagonist. In one embodiment, an antibody of the invention possesses IL-4-blocking activity substantially equivalent to that of L1H11; and possesses IL-13-blocking activity substantially equivalent to that of L1H11. Such activity may be measured in any suitable conventional assay (e.g., as measured in the CD23 expression assay described in Example 2).

20 The DNA sequence of the variable domain of the light chain of L1H11 is presented in SEQ ID NO:3, and the encoded amino acid sequence is presented in SEQ ID NO:4. The DNA sequence for the variable domain of the heavy chain of L1H11 is presented as SEQ ID NO:35, and the encoded amino acid sequence is presented in SEQ ID NO:36. Antibodies of the present invention include, but are not limited to, antibodies that comprise, in their light chain, residues 1 to
25 109 of SEQ ID NO:4; and antibodies that additionally or alternatively comprise, in their heavy chain, residues 1 to 115 of SEQ ID NO:36.

Particular embodiments of antibodies of the present invention comprise, within the variable domain of their light chain, at least one of the complementarity determining regions (CDRs), or hypervariable regions, found in the light chain of L1H11. CDRs of L1H11 are
30 discussed in example 5. Thus, among the antibodies provided herein are those comprising from one to all three of the following sequences in the light chain variable domain: amino acid residues 24-35; residues 51-57; and residues 90-99 of SEQ ID NO:4. Particular antibodies provided herein comprise, within the variable domain of their heavy chain, at least one of the CDRs found in the heavy chain of L1H11. Thus, among the antibodies provided herein are those comprising from
35 one to all three of the following sequences in the heavy chain variable domain: residues 31-35; residues 50-65; and residues 98-104 of SEQ ID NO:36.

In another embodiment, particular antibodies of the invention are selected from the group consisting of L2H4; an antibody that is cross-reactive with L2H4; an antibody that binds to the same epitope as L2H4; an antibody that competes with L2H4 for binding to a cell that expresses
40 human IL-4R; an antibody that possesses a biological activity of L2H4; and an antigen-binding fragment of any of the foregoing antibodies. In one embodiment, the antibody has a binding affinity for human IL-4R that is substantially equivalent to the binding affinity of L2H4 for human IL-4R. Hybridoma cell lines that produce any such antibodies also are provided by the present invention.

5 One example of a biological activity of L2H4 is the ability to function as both an IL-4 antagonist and an IL-13 antagonist. In one embodiment, an antibody of the invention possesses IL-4-blocking activity substantially equivalent to that of L2H4; and possesses IL-13-blocking activity substantially equivalent to that of L2H4. Such activity may be measured in any suitable conventional assay (*e.g.*, as measured in the CD23 expression assay described in Example 2).

10 The DNA sequence of the variable domain of the light chain of L2H4 is presented in SEQ ID NO:5, and the encoded amino acid sequence is presented in SEQ ID NO:6. The DNA sequence for the variable domain of the heavy chain of L2H4 is presented as SEQ ID NO:21, and the encoded amino acid sequence is presented in SEQ ID NO:22. Antibodies of the present invention include, but are not limited to, antibodies that comprise, in their light chain, residues 1 to
15 109 of SEQ ID NO:6; and antibodies that additionally or alternatively comprise, in their heavy chain, residues 1 to 115 of SEQ ID NO:22.

 Particular embodiments of antibodies of the present invention comprise, within the variable domain of their light chain, at least one of the complementarity determining regions (CDRs), or hypervariable regions, found in the light chain of L2H4. CDRs of L2H4 are discussed
20 in example 5. Thus, among the antibodies provided herein are those comprising from one to all three of the following sequences in the light chain variable domain: amino acid residues 24-35; residues 51-57; and residues 90-99 of SEQ ID NO:6. Particular antibodies provided herein comprise, within the variable domain of their heavy chain, at least one of the CDRs found in the heavy chain of L2H4. Thus, among the antibodies provided herein are those comprising from one
25 to all three of the following sequences in the heavy chain variable domain: residues 31-35; residues 50-65; and residues 98-104 of SEQ ID NO:22.

 In another embodiment, particular antibodies of the invention are selected from the group consisting of L2H12; an antibody that is cross-reactive with L2H12; an antibody that binds to the same epitope as L2H12; an antibody that competes with L2H12 for binding to a cell that expresses
30 human IL-4R; an antibody that possesses a biological activity of L2H12; and an antigen-binding fragment of any of the foregoing antibodies. In one embodiment, the antibody has a binding affinity for human IL-4R that is substantially equivalent to the binding affinity of L2H12 for human IL-4R. Hybridoma cell lines that produce any such antibodies also are provided by the present invention.

35 One example of a biological activity of L2H12 is the ability to function as both an IL-4 antagonist and an IL-13 antagonist. In one embodiment, an antibody of the invention possesses IL-4-blocking activity substantially equivalent to that of L2H12; and possesses IL-13-blocking activity substantially equivalent to that of L2H12. Such activity may be measured in any suitable conventional assay (*e.g.*, as measured in the CD23 expression assay described in Example 2).

40 The DNA sequence of the variable domain of the light chain of L2H12 is presented in SEQ ID NO:5, and the encoded amino acid sequence is presented in SEQ ID NO:6. The DNA sequence for the variable domain of the heavy chain of L2H12 is presented as SEQ ID NO:37, and the encoded amino acid sequence is presented in SEQ ID NO:38. Antibodies of the present invention include, but are not limited to, antibodies that comprise, in their light chain, residues 1 to

5 109 of SEQ ID NO:6; and antibodies that additionally or alternatively comprise, in their heavy chain, residues 1 to 115 of SEQ ID NO:38.

Particular embodiments of antibodies of the present invention comprise, within the variable domain of their light chain, at least one of the complementarity determining regions (CDRs), or hypervariable regions, found in the light chain of L2H12. CDRs of L2H12 are
10 discussed in example 5. Thus, among the antibodies provided herein are those comprising from one to all three of the following sequences in the light chain variable domain: amino acid residues 24-35; residues 51-57; and residues 90-99 of SEQ ID NO:6. Particular antibodies provided herein comprise, within the variable domain of their heavy chain, at least one of the CDRs found in the heavy chain of L2H12. Thus, among the antibodies provided herein are those comprising from
15 one to all three of the following sequences in the heavy chain variable domain: residues 31-35; residues 50-65; and residues 98-104 of SEQ ID NO:38.

In another embodiment, particular antibodies of the invention are selected from the group consisting of L2H13; an antibody that is cross-reactive with L2H13; an antibody that binds to the same epitope as L2H13; an antibody that competes with L2H13 for binding to a cell that expresses
20 human IL-4R; an antibody that possesses a biological activity of L2H13; and an antigen-binding fragment of any of the foregoing antibodies. In one embodiment, the antibody has a binding affinity for human IL-4R that is substantially equivalent to the binding affinity of L2H13 for human IL-4R. Hybridoma cell lines that produce any such antibodies also are provided by the present invention.

25 One example of a biological activity of L2H13 is the ability to function as both an IL-4 antagonist and an IL-13 antagonist. In one embodiment, an antibody of the invention possesses IL-4-blocking activity substantially equivalent to that of L2H13; and possesses IL-13-blocking activity substantially equivalent to that of L2H13. Such activity may be measured in any suitable conventional assay (*e.g.*, as measured in the CD23 expression assay described in Example 2).

30 The DNA sequence of the variable domain of the light chain of L2H13 is presented in SEQ ID NO:5, and the encoded amino acid sequence is presented in SEQ ID NO:6. The DNA sequence for the variable domain of the heavy chain of L2H13 is presented as SEQ ID NO:39, and the encoded amino acid sequence is presented in SEQ ID NO:40. Antibodies of the present invention include, but are not limited to, antibodies that comprise, in their light chain, residues 1 to
35 109 of SEQ ID NO:6; and antibodies that additionally or alternatively comprise, in their heavy chain, residues 1 to 115 of SEQ ID NO:40.

Particular embodiments of antibodies of the present invention comprise, within the variable domain of their light chain, at least one of the complementarity determining regions (CDRs), or hypervariable regions, found in the light chain of L2H13. CDRs of L2H13 are
40 discussed in example 5. Thus, among the antibodies provided herein are those comprising from one to all three of the following sequences in the light chain variable domain: amino acid residues 24-35; residues 51-57; and residues 90-99 of SEQ ID NO:6. Particular antibodies provided herein comprise, within the variable domain of their heavy chain, at least one of the CDRs found in the heavy chain of L2H13. Thus, among the antibodies provided herein are those comprising from

5 one to all three of the following sequences in the heavy chain variable domain: residues 31-35; residues 50-65; and residues 98-104 of SEQ ID NO:40.

In another embodiment, particular antibodies of the invention are selected from the group consisting of L2H14; an antibody that is cross-reactive with L2H14; an antibody that binds to the same epitope as L2H14; an antibody that competes with L2H14 for binding to a cell that expresses
10 human IL-4R; an antibody that possesses a biological activity of L2H14; and an antigen-binding fragment of any of the foregoing antibodies. In one embodiment, the antibody has a binding affinity for human IL-4R that is substantially equivalent to the binding affinity of L2H14 for human IL-4R. Hybridoma cell lines that produce any such antibodies also are provided by the present invention.

15 One example of a biological activity of L2H14 is the ability to function as both an IL-4 antagonist and an IL-13 antagonist. In one embodiment, an antibody of the invention possesses IL-4-blocking activity substantially equivalent to that of L2H14; and possesses IL-13-blocking activity substantially equivalent to that of L2H14. Such activity may be measured in any suitable conventional assay (*e.g.*, as measured in the CD23 expression assay described in Example 2).

20 The DNA sequence of the variable domain of the light chain of L2H14 is presented in SEQ ID NO:5, and the encoded amino acid sequence is presented in SEQ ID NO:6. The DNA sequence for the variable domain of the heavy chain of L2H14 is presented as SEQ ID NO:41, and the encoded amino acid sequence is presented in SEQ ID NO:42. Antibodies of the present invention include, but are not limited to, antibodies that comprise, in their light chain, residues 1 to
25 109 of SEQ ID NO:6; and antibodies that additionally or alternatively comprise, in their heavy chain, residues 1 to 115 of SEQ ID NO:42.

Particular embodiments of antibodies of the present invention comprise, within the variable domain of their light chain, at least one of the complementarity determining regions (CDRs), or hypervariable regions, found in the light chain of L2H14. CDRs of L2H14 are
30 discussed in example 5. Thus, among the antibodies provided herein are those comprising from one to all three of the following sequences in the light chain variable domain: amino acid residues 24-35; residues 51-57; and residues 90-99 of SEQ ID NO:6. Particular antibodies provided herein comprise, within the variable domain of their heavy chain, at least one of the CDRs found in the heavy chain of L2H14. Thus, among the antibodies provided herein are those comprising from
35 one to all three of the following sequences in the heavy chain variable domain: residues 31-35; residues 50-65; and residues 98-104 of SEQ ID NO:42.

In another embodiment, particular antibodies of the invention are selected from the group consisting of L6H1; an antibody that is cross-reactive with L6H1; an antibody that binds to the same epitope as L6H1; an antibody that competes with L6H1 for binding to a cell that expresses
40 human IL-4R; an antibody that possesses a biological activity of L6H1; and an antigen-binding fragment of any of the foregoing antibodies. In one embodiment, the antibody has a binding affinity for human IL-4R that is substantially equivalent to the binding affinity of L6H1 for human IL-4R. Hybridoma cell lines that produce any such antibodies also are provided by the present invention.

5 One example of a biological activity of L6H1 is the ability to function as both an IL-4 antagonist and an IL-13 antagonist. In one embodiment, an antibody of the invention possesses IL-4-blocking activity substantially equivalent to that of L6H1; and possesses IL-13-blocking activity substantially equivalent to that of L6H1. Such activity may be measured in any suitable conventional assay (e.g., as measured in the CD23 expression assay described in Example 2).

10 The DNA sequence of the variable domain of the light chain of L6H1 is presented in SEQ ID NO:13, and the encoded amino acid sequence is presented in SEQ ID NO:14. The DNA sequence for the variable domain of the heavy chain of L6H1 is presented as SEQ ID NO:15, and the encoded amino acid sequence is presented in SEQ ID NO:16. Antibodies of the present invention include, but are not limited to, antibodies that comprise, in their light chain, residues 1 to
15 109 of SEQ ID NO:14; and antibodies that additionally or alternatively comprise, in their heavy chain, residues 1 to 115 of SEQ ID NO:16.

 Particular embodiments of antibodies of the present invention comprise, within the variable domain of their light chain, at least one of the complementarity determining regions (CDRs), or hypervariable regions, found in the light chain of L6H1. CDRs of L6H1 are discussed
20 in example 5. Thus, among the antibodies provided herein are those comprising from one to all three of the following sequences in the light chain variable domain: amino acid residues 24-35; residues 51-57; and residues 90-99 of SEQ ID NO:14. Particular antibodies provided herein comprise, within the variable domain of their heavy chain, at least one of the CDRs found in the heavy chain of L6H1. Thus, among the antibodies provided herein are those comprising from one
25 to all three of the following sequences in the heavy chain variable domain: residues 31-35; residues 50-65; and residues 98-104 of SEQ ID NO:16.

 In another embodiment, particular antibodies of the invention are selected from the group consisting of L2H2; an antibody that is cross-reactive with L2H2; an antibody that binds to the same epitope as L2H2; an antibody that competes with L2H2 for binding to a cell that expresses
30 human IL-4R; an antibody that possesses a biological activity of L2H2; and an antigen-binding fragment of any of the foregoing antibodies. In one embodiment, the antibody has a binding affinity for human IL-4R that is substantially equivalent to the binding affinity of L2H2 for human IL-4R. Hybridoma cell lines that produce any such antibodies also are provided by the present invention.

35 One example of a biological activity of L2H2 is the ability to function as both an IL-4 antagonist and an IL-13 antagonist. In one embodiment, an antibody of the invention possesses IL-4-blocking activity substantially equivalent to that of L2H2; and possesses IL-13-blocking activity substantially equivalent to that of L2H2. Such activity may be measured in any suitable conventional assay (e.g., as measured in the CD23 expression assay described in Example 2).

40 The DNA sequence of the variable domain of the light chain of L2H2 is presented in SEQ ID NO:5, and the encoded amino acid sequence is presented in SEQ ID NO:6. The DNA sequence for the variable domain of the heavy chain of L2H2 is presented as SEQ ID NO:17, and the encoded amino acid sequence is presented in SEQ ID NO:18. Antibodies of the present invention include, but are not limited to, antibodies that comprise, in their light chain, residues 1 to

5 109 of SEQ ID NO:6; and antibodies that additionally or alternatively comprise, in their heavy chain, residues 1 to 115 of SEQ ID NO:18.

Particular embodiments of antibodies of the present invention comprise, within the variable domain of their light chain, at least one of the complementarity determining regions (CDRs), or hypervariable regions, found in the light chain of L2H2. CDRs of L2H2 are discussed
10 in example 5. Thus, among the antibodies provided herein are those comprising from one to all three of the following sequences in the light chain variable domain: amino acid residues 24-35; residues 51-57; and residues 90-99 of SEQ ID NO:6. Particular antibodies provided herein comprise, within the variable domain of their heavy chain, at least one of the CDRs found in the heavy chain of L2H2. Thus, among the antibodies provided herein are those comprising from one
15 to all three of the following sequences in the heavy chain variable domain: residues 31-35; residues 50-65; and residues 98-104 of SEQ ID NO:18.

In another embodiment, particular antibodies of the invention are selected from the group consisting of L2H3; an antibody that is cross-reactive with L2H3; an antibody that binds to the same epitope as L2H3; an antibody that competes with L2H3 for binding to a cell that expresses
20 human IL-4R; an antibody that possesses a biological activity of L2H3; and an antigen-binding fragment of any of the foregoing antibodies. In one embodiment, the antibody has a binding affinity for human IL-4R that is substantially equivalent to the binding affinity of L2H3 for human IL-4R. Hybridoma cell lines that produce any such antibodies also are provided by the present invention.

25 One example of a biological activity of L2H3 is the ability to function as both an IL-4 antagonist and an IL-13 antagonist. In one embodiment, an antibody of the invention possesses IL-4-blocking activity substantially equivalent to that of L2H3; and possesses IL-13-blocking activity substantially equivalent to that of L2H3. Such activity may be measured in any suitable conventional assay (e.g., as measured in the CD23 expression assay described in Example 2).

30 The DNA sequence of the variable domain of the light chain of L2H3 is presented in SEQ ID NO:5, and the encoded amino acid sequence is presented in SEQ ID NO:6. The DNA sequence for the variable domain of the heavy chain of L2H3 is presented as SEQ ID NO:19, and the encoded amino acid sequence is presented in SEQ ID NO:20. Antibodies of the present invention include, but are not limited to, antibodies that comprise, in their light chain, residues 1 to
35 109 of SEQ ID NO:6; and antibodies that additionally or alternatively comprise, in their heavy chain, residues 1 to 115 of SEQ ID NO:20.

Particular embodiments of antibodies of the present invention comprise, within the variable domain of their light chain, at least one of the complementarity determining regions (CDRs), or hypervariable regions, found in the light chain of L2H3. CDRs of L2H3 are discussed
40 in example 5. Thus, among the antibodies provided herein are those comprising from one to all three of the following sequences in the light chain variable domain: amino acid residues 24-35; residues 51-57; and residues 90-99 of SEQ ID NO:6. Particular antibodies provided herein comprise, within the variable domain of their heavy chain, at least one of the CDRs found in the heavy chain of L2H3. Thus, among the antibodies provided herein are those comprising from one

5 to all three of the following sequences in the heavy chain variable domain: residues 31-35; residues 50-65; and residues 98-104 of SEQ ID NO:20.

In another embodiment, particular antibodies of the invention are selected from the group consisting of L2H6; an antibody that is cross-reactive with L2H6; an antibody that binds to the same epitope as L2H6; an antibody that competes with L2H6 for binding to a cell that expresses
10 human IL-4R; an antibody that possesses a biological activity of L2H6; and an antigen-binding fragment of any of the foregoing antibodies. In one embodiment, the antibody has a binding affinity for human IL-4R that is substantially equivalent to the binding affinity of L2H6 for human IL-4R. Hybridoma cell lines that produce any such antibodies also are provided by the present invention.

15 One example of a biological activity of L2H6 is the ability to function as both an IL-4 antagonist and an IL-13 antagonist. In one embodiment, an antibody of the invention possesses IL-4-blocking activity substantially equivalent to that of L2H6; and possesses IL-13-blocking activity substantially equivalent to that of L2H6. Such activity may be measured in any suitable conventional assay (*e.g.*, as measured in the CD23 expression assay described in Example 2).

20 The DNA sequence of the variable domain of the light chain of L2H6 is presented in SEQ ID NO:5, and the encoded amino acid sequence is presented in SEQ ID NO:6. The DNA sequence for the variable domain of the heavy chain of L2H6 is presented as SEQ ID NO:25, and the encoded amino acid sequence is presented in SEQ ID NO:26. Antibodies of the present invention include, but are not limited to, antibodies that comprise, in their light chain, residues 1 to
25 109 of SEQ ID NO:6; and antibodies that additionally or alternatively comprise, in their heavy chain, residues 1 to 115 of SEQ ID NO:26.

Particular embodiments of antibodies of the present invention comprise, within the variable domain of their light chain, at least one of the complementarity determining regions (CDRs), or hypervariable regions, found in the light chain of L2H6. CDRs of L2H6 are discussed
30 in example 5. Thus, among the antibodies provided herein are those comprising from one to all three of the following sequences in the light chain variable domain: amino acid residues 24-35; residues 51-57; and residues 90-99 of SEQ ID NO:6. Particular antibodies provided herein comprise, within the variable domain of their heavy chain, at least one of the CDRs found in the heavy chain of L2H6. Thus, among the antibodies provided herein are those comprising from one
35 to all three of the following sequences in the heavy chain variable domain: residues 31-35; residues 50-65; and residues 98-104 of SEQ ID NO:26.

In another embodiment, particular antibodies of the invention are selected from the group consisting of L2H7; an antibody that is cross-reactive with L2H7; an antibody that binds to the same epitope as L2H7; an antibody that competes with L2H7 for binding to a cell that expresses
40 human IL-4R; an antibody that possesses a biological activity of L2H7; and an antigen-binding fragment of any of the foregoing antibodies. In one embodiment, the antibody has a binding affinity for human IL-4R that is substantially equivalent to the binding affinity of L2H7 for human IL-4R. Hybridoma cell lines that produce any such antibodies also are provided by the present invention.

5 One example of a biological activity of L2H7 is the ability to function as both an IL-4 antagonist and an IL-13 antagonist. In one embodiment, an antibody of the invention possesses IL-4-blocking activity substantially equivalent to that of L2H7; and possesses IL-13-blocking activity substantially equivalent to that of L2H7. Such activity may be measured in any suitable conventional assay (e.g., as measured in the CD23 expression assay described in Example 2).

10 The DNA sequence of the variable domain of the light chain of L2H7 is presented in SEQ ID NO:5, and the encoded amino acid sequence is presented in SEQ ID NO:6. The DNA sequence for the variable domain of the heavy chain of L2H7 is presented as SEQ ID NO:27, and the encoded amino acid sequence is presented in SEQ ID NO:28. Antibodies of the present invention include, but are not limited to, antibodies that comprise, in their light chain, residues 1 to
15 109 of SEQ ID NO:6; and antibodies that additionally or alternatively comprise, in their heavy chain, residues 1 to 115 of SEQ ID NO:28.

 Particular embodiments of antibodies of the present invention comprise, within the variable domain of their light chain, at least one of the complementarity determining regions (CDRs), or hypervariable regions, found in the light chain of L2H7. CDRs of L2H7 are discussed
20 in example 5. Thus, among the antibodies provided herein are those comprising from one to all three of the following sequences in the light chain variable domain: amino acid residues 24-35; residues 51-57; and residues 90-99 of SEQ ID NO:6. Particular antibodies provided herein comprise, within the variable domain of their heavy chain, at least one of the CDRs found in the heavy chain of L2H7. Thus, among the antibodies provided herein are those comprising from one
25 to all three of the following sequences in the heavy chain variable domain: residues 31-35; residues 50-65; and residues 98-104 of SEQ ID NO:28.

 In another embodiment, particular antibodies of the invention are selected from the group consisting of L2H8; an antibody that is cross-reactive with L2H8; an antibody that binds to the same epitope as L2H8; an antibody that competes with L2H8 for binding to a cell that expresses
30 human IL-4R; an antibody that possesses a biological activity of L2H8; and an antigen-binding fragment of any of the foregoing antibodies. In one embodiment, the antibody has a binding affinity for human IL-4R that is substantially equivalent to the binding affinity of L2H8 for human IL-4R. Hybridoma cell lines that produce any such antibodies also are provided by the present invention.

35 One example of a biological activity of L2H8 is the ability to function as both an IL-4 antagonist and an IL-13 antagonist. In one embodiment, an antibody of the invention possesses IL-4-blocking activity substantially equivalent to that of L2H8; and possesses IL-13-blocking activity substantially equivalent to that of L2H8. Such activity may be measured in any suitable conventional assay (e.g., as measured in the CD23 expression assay described in Example 2).

40 The DNA sequence of the variable domain of the light chain of L2H8 is presented in SEQ ID NO:5, and the encoded amino acid sequence is presented in SEQ ID NO:6. The DNA sequence for the variable domain of the heavy chain of L2H8 is presented as SEQ ID NO:29, and the encoded amino acid sequence is presented in SEQ ID NO:30. Antibodies of the present invention include, but are not limited to, antibodies that comprise, in their light chain, residues 1 to

5 109 of SEQ ID NO:6; and antibodies that additionally or alternatively comprise, in their heavy chain, residues 1 to 115 of SEQ ID NO:30.

Particular embodiments of antibodies of the present invention comprise, within the variable domain of their light chain, at least one of the complementarity determining regions (CDRs), or hypervariable regions, found in the light chain of L2H8. CDRs of L2H8 are discussed
10 in example 5. Thus, among the antibodies provided herein are those comprising from one to all three of the following sequences in the light chain variable domain: amino acid residues 24-35; residues 51-57; and residues 90-99 of SEQ ID NO:6. Particular antibodies provided herein comprise, within the variable domain of their heavy chain, at least one of the CDRs found in the heavy chain of L2H8. Thus, among the antibodies provided herein are those comprising from one
15 to all three of the following sequences in the heavy chain variable domain: residues 31-35; residues 50-65; and residues 98-104 of SEQ ID NO:30.

In another embodiment, particular antibodies of the invention are selected from the group consisting of L2H9; an antibody that is cross-reactive with L2H9; an antibody that binds to the same epitope as L2H9; an antibody that competes with L2H9 for binding to a cell that expresses
20 human IL-4R; an antibody that possesses a biological activity of L2H9; and an antigen-binding fragment of any of the foregoing antibodies. In one embodiment, the antibody has a binding affinity for human IL-4R that is substantially equivalent to the binding affinity of L2H9 for human IL-4R. Hybridoma cell lines that produce any such antibodies also are provided by the present invention.

25 One example of a biological activity of L2H9 is the ability to function as both an IL-4 antagonist and an IL-13 antagonist. In one embodiment, an antibody of the invention possesses IL-4-blocking activity substantially equivalent to that of L2H9; and possesses IL-13-blocking activity substantially equivalent to that of L2H9. Such activity may be measured in any suitable conventional assay (e.g., as measured in the CD23 expression assay described in Example 2).

30 The DNA sequence of the variable domain of the light chain of L2H9 is presented in SEQ ID NO:5, and the encoded amino acid sequence is presented in SEQ ID NO:6. The DNA sequence for the variable domain of the heavy chain of L2H9 is presented as SEQ ID NO:31, and the encoded amino acid sequence is presented in SEQ ID NO:32. Antibodies of the present invention include, but are not limited to, antibodies that comprise, in their light chain, residues 1 to
35 109 of SEQ ID NO:6; and antibodies that additionally or alternatively comprise, in their heavy chain, residues 1 to 115 of SEQ ID NO:32.

Particular embodiments of antibodies of the present invention comprise, within the variable domain of their light chain, at least one of the complementarity determining regions (CDRs), or hypervariable regions, found in the light chain of L2H9. CDRs of L2H9 are discussed
40 in example 5. Thus, among the antibodies provided herein are those comprising from one to all three of the following sequences in the light chain variable domain: amino acid residues 24-35; residues 51-57; and residues 90-99 of SEQ ID NO:6. Particular antibodies provided herein comprise, within the variable domain of their heavy chain, at least one of the CDRs found in the heavy chain of L2H9. Thus, among the antibodies provided herein are those comprising from one

5 to all three of the following sequences in the heavy chain variable domain: residues 31-35; residues 50-65; and residues 98-104 of SEQ ID NO:32.

In another embodiment, particular antibodies of the invention are selected from the group consisting of L2H10; an antibody that is cross-reactive with L2H10; an antibody that binds to the same epitope as L2H10; an antibody that competes with L2H10 for binding to a cell that expresses
10 human IL-4R; an antibody that possesses a biological activity of L2H10; and an antigen-binding fragment of any of the foregoing antibodies. In one embodiment, the antibody has a binding affinity for human IL-4R that is substantially equivalent to the binding affinity of L2H10 for human IL-4R. Hybridoma cell lines that produce any such antibodies also are provided by the present invention.

15 One example of a biological activity of L2H10 is the ability to function as both an IL-4 antagonist and an IL-13 antagonist. In one embodiment, an antibody of the invention possesses IL-4-blocking activity substantially equivalent to that of L2H10; and possesses IL-13-blocking activity substantially equivalent to that of L2H10. Such activity may be measured in any suitable conventional assay (*e.g.*, as measured in the CD23 expression assay described in Example 2).

20 The DNA sequence of the variable domain of the light chain of L2H10 is presented in SEQ ID NO:5, and the encoded amino acid sequence is presented in SEQ ID NO:6. The DNA sequence for the variable domain of the heavy chain of L2H10 is presented as SEQ ID NO:33, and the encoded amino acid sequence is presented in SEQ ID NO:34. Antibodies of the present invention include, but are not limited to, antibodies that comprise, in their light chain, residues 1 to
25 109 of SEQ ID NO:6; and antibodies that additionally or alternatively comprise, in their heavy chain, SEQ ID NO:34.

Particular embodiments of antibodies of the present invention comprise, within the variable domain of their light chain, at least one of the complementarity determining regions (CDRs), or hypervariable regions, found in the light chain of L2H10. CDRs of L2H10 are
30 discussed in example 5. Thus, among the antibodies provided herein are those comprising from one to all three of the following sequences in the light chain variable domain: amino acid residues 24-35; residues 51-57; and residues 90-99 of SEQ ID NO:6. Particular antibodies provided herein comprise, within the variable domain of their heavy chain, at least one of the CDRs found in the heavy chain of L2H10. Thus, among the antibodies provided herein are those comprising from
35 one to all three of the following sequences in the heavy chain variable domain: residues 31-35; residues 50-65; and residues 98-104 of SEQ ID NO:34.

In another embodiment, particular antibodies of the invention are selected from the group consisting of L2H11; an antibody that is cross-reactive with L2H11; an antibody that binds to the same epitope as L2H11; an antibody that competes with L2H11 for binding to a cell that expresses
40 human IL-4R; an antibody that possesses a biological activity of L2H11; and an antigen-binding fragment of any of the foregoing antibodies. In one embodiment, the antibody has a binding affinity for human IL-4R that is substantially equivalent to the binding affinity of L2H11 for human IL-4R. Hybridoma cell lines that produce any such antibodies also are provided by the present invention.

5 One example of a biological activity of L2H11 is the ability to function as both an IL-4 antagonist and an IL-13 antagonist. In one embodiment, an antibody of the invention possesses IL-4-blocking activity substantially equivalent to that of L2H11; and possesses IL-13-blocking activity substantially equivalent to that of L2H11. Such activity may be measured in any suitable conventional assay (e.g., as measured in the CD23 expression assay described in Example 2).

10 The DNA sequence of the variable domain of the light chain of L2H11 is presented in SEQ ID NO:5, and the encoded amino acid sequence is presented in SEQ ID NO:6. The DNA sequence for the variable domain of the heavy chain of L2H11 is presented as SEQ ID NO:35, and the encoded amino acid sequence is presented in SEQ ID NO:36. Antibodies of the present invention include, but are not limited to, antibodies that comprise, in their light chain, residues 1 to
15 109 of SEQ ID NO:6; and antibodies that additionally or alternatively comprise, in their heavy chain, residues 1 to 115 of SEQ ID NO:36.

Particular embodiments of antibodies of the present invention comprise, within the variable domain of their light chain, at least one of the complementarity determining regions (CDRs), or hypervariable regions, found in the light chain of L2H11. CDRs of L2H11 are
20 discussed in example 5. Thus, among the antibodies provided herein are those comprising from one to all three of the following sequences in the light chain variable domain: amino acid residues 24-35; residues 51-57; and residues 90-99 of SEQ ID NO:6. Particular antibodies provided herein comprise, within the variable domain of their heavy chain, at least one of the CDRs found in the heavy chain of L2H11. Thus, among the antibodies provided herein are those comprising from
25 one to all three of the following sequences in the heavy chain variable domain: residues 31-35; residues 50-65; and residues 98-104 of SEQ ID NO:36.

Nucleic acids

30 In one aspect, the present invention provides isolated nucleic acid molecules. The nucleic acids comprise, for example, polynucleotides that encode an antibody of the invention, or a fragment, derivative, mutein, or variant thereof, polynucleotides sufficient for use as hybridization probes, PCR primers or sequencing primers for identifying, analyzing, mutating or amplifying a polynucleotide encoding a polypeptide, anti-sense nucleic acids for inhibiting expression of a polynucleotide, and complementary sequences of the foregoing. The nucleic acids can be any
35 length. They can be, for example, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 75, 100, 125, 150, 175, 200, 250, 300, 350, 400, 450, 500, 750, 1,000, 1,500, 3,000, 5,000 or more nucleotides in length, and/or can comprise one or more additional sequences, for example, regulatory sequences, and/or be part of a larger nucleic acid, for example, a vector. The nucleic acids can be single-stranded or double-stranded and can comprise RNA and/or DNA nucleotides, and artificial
40 variants thereof (e.g., peptide nucleic acids).

DNA encoding antibody polypeptides (e.g., heavy or light chain, variable domain only or full length) may be isolated from B-cells of mice that have been immunized with IL-4R. The DNA may be isolated by conventional procedures such as polymerase chain reaction (PCR). Phage display is another example of a known technique whereby derivatives of antibodies may be
45 prepared. In one approach, polypeptides that are components of an antibody of interest are

5 expressed in any suitable recombinant expression system, and the expressed polypeptides are allowed to assemble to form antibody molecules.

Figure 2 provides nucleic acid sequences encoding the variable regions of the heavy and light chain variable regions shown in Figure 3. Due to the degeneracy of the genetic code, each of the polypeptide sequences in Figure 3 also is encoded by a large number of other nucleic acid sequences. The present invention provides each degenerate nucleotide sequence encoding each antibody of the invention.

The invention further provides nucleic acids that hybridize to other nucleic acids (*e.g.*, nucleic acids comprising a nucleotide sequence of Figure 2) under particular hybridization conditions. Methods for hybridizing nucleic acids are well-known in the art. See, *e.g.*, Current Protocols in Molecular Biology, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6. As defined herein, a moderately stringent hybridization condition uses a prewashing solution containing 5X sodium chloride/sodium citrate (SSC), 0.5% SDS, 1.0 mM EDTA (pH 8.0), hybridization buffer of about 50% formamide, 6X SSC, and a hybridization temperature of 55° C (or other similar hybridization solutions, such as one containing about 50% formamide, with a hybridization temperature of 42° C), and washing conditions of 60° C, in 0.5X SSC, 0.1% SDS. A stringent hybridization condition 6X SSC at 45° C, followed by one or more washes in 0.1X SSC, 0.2% SDS at 68° C.

Furthermore, one of skill in the art can manipulate the hybridization and/or washing conditions to increase or decrease the stringency of hybridization such that nucleic acids comprising nucleotide sequences that are at least 65, 70, 75, 80, 85, 90, 95, 98 or 99% identical to each other typically remain hybridized to each other. The basic parameters affecting the choice of hybridization conditions and guidance for devising suitable conditions are set forth by Sambrook, Fritsch, and Maniatis (1989, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., chapters 9 and 11; and Current Protocols in Molecular Biology, 1995, Ausubel *et al.*, eds., John Wiley & Sons, Inc., sections 2.10 and 6.3-6.4), and can be readily determined by those having ordinary skill in the art based on, for example, the length and/or base composition of the DNA.

Changes can be introduced by mutation into a nucleic acid, thereby leading to changes in the amino acid sequence of a polypeptide (*e.g.*, an antibody) that it encodes. Mutations can be introduced by standard techniques, such as site-directed mutagenesis and PCR-mediated mutagenesis. Preferably, conservative amino acid substitutions are made at one or more predicted non-essential amino acid residues. Alternatively, mutations can be introduced randomly along all or part of the coding sequence, such as by saturation mutagenesis, and the resultant mutants can be screened for biological activity to identify mutants that retain activity (*e.g.*, binding to IL-4 receptor). Following mutagenesis, the encoded protein can be expressed recombinantly and the activity of the protein can be determined. The mutations can be introduced into a nucleic acid without significantly altering the biological activity of a peptide that it encodes. For example, one can make nucleotide substitutions leading to amino acid substitutions at non-essential amino acid residues. In one embodiment, a nucleotide sequence provided in Figure 2, or a desired fragment, variant, or derivative thereof, is mutated such that it encodes an amino acid sequence comprising one or more deletions or substitutions of amino acid residues that are shown in Figure

5 3 to be residues where two or more sequences differ. In another embodiment, the mutagenesis inserts an amino acid adjacent to one or more amino acid residues shown in Figure 3 to be residues where two or more sequences differ. Alternatively, mutations can be introduced into a nucleic acid that selectively change the biological activity (*e.g.*, binding of IL-4 receptor, inhibiting IL-4 and/or IL-13, *etc.*) of a polypeptide that it encodes. For example, the mutation can
10 quantitatively or qualitatively change the biological activity. Examples of quantitative changes include increasing, reducing or eliminating the activity. Examples of qualitative changes include changing the antigen specificity of an antibody.

In another aspect, the present invention provides nucleic acid molecules that are suitable for use as primers or hybridization probes for the detection of nucleic acid sequences of the
15 invention. A nucleic acid molecule of the invention can comprise only a portion of a nucleic acid sequence encoding a full length polypeptide of the invention, for example, a fragment that can be used as a probe or primer or a fragment encoding an active portion (*e.g.*, an IL-4 receptor binding portion) of a polypeptide of the invention.

Probes based on the sequence of a nucleic acid of the invention can be used to detect the
20 nucleic acid or similar nucleic acids, for example, transcripts encoding a polypeptide of the invention. The probe can comprise a label group, *e.g.*, a radioisotope, a fluorescent compound, an enzyme, or an enzyme co-factor. Such probes can be used identify a cell that expresses the polypeptide.

In another aspect, the present invention provides vectors comprising a nucleic acid
25 encoding a polypeptide of the invention or a portion thereof. Examples of vectors include, but are not limited to, plasmids, viral vectors, non-episomal mammalian vectors and expression vectors, for example, recombinant expression vectors.

The recombinant expression vectors of the invention can comprise a nucleic acid of the invention in a form suitable for expression of the nucleic acid in a host cell. The recombinant
30 expression vectors include one or more regulatory sequences, selected on the basis of the host cells to be used for expression, which is operably linked to the nucleic acid sequence to be expressed. Regulatory sequences include those that direct constitutive expression of a nucleotide sequence in many types of host cells (*e.g.*, SV40 early gene enhancer, Rous sarcoma virus promoter and cytomegalovirus promoter), those that direct expression of the nucleotide sequence only in certain host cells (*e.g.*, tissue-specific regulatory sequences, see Voss *et al.*, 1986, Trends
35 Biochem. Sci. 11:287, Maniatis *et al.*, 1987, Science 236:1237, incorporated by reference herein in their entireties), and those that direct inducible expression of a nucleotide sequence in response to particular treatment or condition (*e.g.*, the metallothionin promoter in mammalian cells and the tet-responsive and/or streptomycin responsive promoter in both prokaryotic and eukaryotic systems
40 (see *id.*). It will be appreciated by those skilled in the art that the design of the expression vector can depend on such factors as the choice of the host cell to be transformed, the level of expression of protein desired, *etc.* The expression vectors of the invention can be introduced into host cells to thereby produce proteins or peptides, including fusion proteins or peptides, encoded by nucleic acids as described herein.

5 In another aspect, the present invention provides host cells into which a recombinant expression vector of the invention has been introduced. A host cell can be any prokaryotic cell (for example, *E. coli*) or eukaryotic cell (for example, yeast, insect, or mammalian cells (*e.g.*, CHO cells)). Vector DNA can be introduced into prokaryotic or eukaryotic cells via conventional transformation or transfection techniques. For stable transfection of mammalian cells, it is known
10 that, depending upon the expression vector and transfection technique used, only a small fraction of cells may integrate the foreign DNA into their genome. In order to identify and select these integrants, a gene that encodes a selectable marker (*e.g.*, for resistance to antibiotics) is generally introduced into the host cells along with the gene of interest. Preferred selectable markers include those which confer resistance to drugs, such as G418, hygromycin and methotrexate. Cells stably
15 transfected with the introduced nucleic acid can be identified by drug selection (*e.g.*, cells that have incorporated the selectable marker gene will survive, while the other cells die), among other methods.

Derivatives, fragments, and muteins of antibodies

20 Derivatives of antibodies directed against IL-4R may be prepared, and screened for desired properties, by any of a number of known techniques. Certain of the techniques involve isolating DNA encoding a polypeptide chain (or portion thereof) of an antibody of interest, and manipulating the DNA through recombinant DNA technology. The DNA may be fused to another DNA of interest, or altered (*e.g.*, by mutagenesis or other conventional techniques) to add, delete,
25 or substitute one or more amino acid residues, for example.

In one aspect, the present invention provides antigen-binding fragments of the antibodies of the invention. Such fragments can consist entirely of antibody-derived sequences or can comprise additional sequences. Examples of antigen-binding fragments include Fab and F(ab')₂. Other examples are provided in Lunde *et al.*, 2002, *Biochem. Soc. Trans.* 30:500-06.

30 Single chain antibodies may be formed by linking heavy and light chain variable domain (Fv region) fragments via an amino acid bridge (short peptide linker), resulting in a single polypeptide chain. Such single-chain Fvs (scFvs) have been prepared by fusing DNA encoding a peptide linker between DNAs encoding the two variable domain polypeptides (V_L and V_H). The resulting polypeptides can fold back on themselves to form antigen-binding monomers, or they
35 can form multimers (*e.g.*, dimers, trimers, or tetramers), depending on the length of a flexible linker between the two variable domains (Kortt *et al.*, 1997, *Prot. Eng.* 10:423; Kortt *et al.*, 2001, *Biomol. Eng.* 18:95-108). By combining different V_L and V_H-comprising polypeptides, one can form multimeric scFvs that bind to different epitopes (Kriangkum *et al.*, 2001, *Biomol. Eng.* 18:31-40). Techniques developed for the production of single chain antibodies include those described
40 in U.S. Patent No. 4,946,778; Bird, 1988, *Science* 242:423; Huston *et al.*, 1988, *Proc. Natl. Acad. Sci. USA* 85:5879; Ward *et al.*, 1989, *Nature* 334:544; de Graaf *et al.*, 2002, *Methods Mol Biol.* 178:379-87. Single chain antibodies derived from antibodies provided herein (including but not limited to scFvs comprising the variable domain combinations L2H1, L3H1, L4H1, L5H1, L1H2, L1H3, L1H4, L1H5, L1H6, L1H7, L1H8, L1H9, L1H10, L1H11, L2H4, L2H12, L2H13, L2H14,

5 L6H1, L2H2, L2H3, L2H6, L2H7, L2H8, L2H9, L2H10, and L2H11) are encompassed by the present invention.

Techniques are known for deriving an antibody of a different subclass or isotype from an antibody of interest, *i.e.*, subclass switching. Thus, IgG antibodies may be derived from an IgM antibody, for example, and *vice versa*. Such techniques allow the preparation of new antibodies that possess the antigen-binding properties of a given antibody (the parent antibody), but also exhibit biological properties associated with an antibody isotype or subclass different from that of the parent antibody. Recombinant DNA techniques may be employed. Cloned DNA encoding particular antibody polypeptides may be employed in such procedures, *e.g.*, DNA encoding the constant domain of an antibody of the desired isotype. See also Lantto *et al.*, 2002, Methods Mol. Biol. 178:303-16.

Accordingly, the antibodies of the present invention include those comprising, for example, the variable domain combinations L2H1, L3H1, L4H1, L5H1, L1H2, L1H3, L1H4, L1H5, L1H6, L1H7, L1H8, L1H9, L1H10, L1H11, L2H4, L2H12, L2H13, L2H14, L6H1, L2H2, L2H3, L2H6, L2H7, L2H8, L2H9, L2H10, and L2H11, having a desired isotype (for example, IgA, IgG1, IgG2, IgG3, IgG4, IgM, IgE, and IgD) as well as Fab or F(ab')₂ fragments thereof. Moreover, if an IgG4 is desired, it may also be desired to introduce a point mutation (CPSCP → CPPCP) in the hinge region as described in Bloom *et al.*, 1997, Protein Science 6:407, incorporated by reference herein) to alleviate a tendency to form intra-H chain disulfide bonds that can lead to heterogeneity in the IgG4 antibodies.

Moreover, techniques for deriving antibodies having different properties (*i.e.*, varying affinities for the antigen to which they bind) are also known. One such technique, referred to as chain shuffling, involves displaying immunoglobulin variable domain gene repertoires on the surface of filamentous bacteriophage, often referred to as phage display. Chain shuffling has been used to prepare high affinity antibodies to the hapten 2-phenyloxazol-5-one, as described by Marks *et al.*, 1992, BioTechnology, 10:779.

Molecular evolution of the complementarity determining regions (CDRs) in the center of the antibody binding site has also been used to isolate antibodies with increased affinity, for example, antibodies having increased affinity for c-erbB-2, as described by Schier *et al.*, 1996, J. Mol. Biol. 263:551. Accordingly, such techniques are useful in preparing antibodies to IL-4R.

In particular embodiments, antibodies of the present invention have a binding affinity (K_a) for IL-4R of at least 1×10^8 . In other embodiments, the antibodies exhibit a K_a of at least 1×10^9 , at least 1×10^{10} or at least 1×10^{11} .

The present invention further includes multi-specific antibodies, for example, bispecific antibodies, *e.g.*, two different epitopes of IL-4R, or an epitope of IL-4R and an epitope of IL-13R, via two different antigen binding sites or regions. Moreover, bispecific antibodies as disclosed herein can comprise an antigen binding site from one of the herein described antibodies and a second antigen binding region from another of the herein described antibodies. Alternatively, a bispecific antibody may comprise an antigen binding site from one of the herein described

5 antibodies and a second antigen binding site from another IL-4R antibody that is known in the art (or one that can be prepared by known methods).

Numerous methods of preparing bispecific antibodies are known in the art, and discussed in US Patent Application 09/839,632, filed April 20, 2001 (incorporated by reference herein). Such methods include the use of hybrid-hybridomas as described by Milstein *et al.*, 1983, Nature
10 305:537, and others (U.S. Patent 4,474,893, U.S. Patent 6,106,833), and chemical coupling of antibody fragments (Brennan *et al.*, 1985, Science 229:81; Glennie *et al.*, 1987, J. Immunol. 139:2367; U.S. Patent 6,010,902). Moreover, bispecific antibodies can be produced via recombinant means, for example by using leucine zipper moieties (*i.e.*, from the *Fos* and *Jun* proteins, which preferentially form heterodimers; Kostelny *et al.*, 1992, J. Immunol. 148:1547) or
15 other lock and key interactive domain structures as described in U.S. Patent 5,582,996. Additional useful techniques include those described in Kortt *et al.*, 1997, *supra*; U.S. Patent 5,959,083; and U.S. Patent 5,807,706.

In another aspect, the present invention provides a derivative of an antibody. The derivatized antibody can comprise any molecule or substance that imparts a desired property to
20 the antibody, such as increased half-life in a particular use. The derivatized antibody can comprise, for example, a detectable (or labeling) moiety (*e.g.*, a radioactive, colorimetric, antigenic or enzymatic molecule, a detectable bead (such as a magnetic or gold bead), or a molecule that binds to another molecule (*e.g.*, biotin)), a therapeutic or diagnostic moiety (*e.g.*, a radioactive, cytotoxic, or pharmaceutically active moiety), or a molecule that increases the suitability of the
25 antibody for a particular use (*e.g.*, administration to a subject, such as a human subject, or other *in vivo* or *in vitro* uses). Examples of molecules that can be used to derivatize an antibody are albumin (*e.g.*, human serum albumin) and polyethylene glycol (PEG). Albumin-linked and PEGylated derivatives of antibodies can be prepared using techniques well known in the art. In one embodiment, the antibody is conjugated or otherwise linked to transthyretin (TTR) or a TTR
30 variant. The TTR or TTR variant can be chemically modified with, for example, a chemical selected from the group consisting of dextran, poly(n-vinyl pyrrolidone), polyethylene glycols, propylene glycol homopolymers, polypropylene oxide/ethylene oxide co-polymers, polyoxyethylated polyols and polyvinyl alcohols. US Pat. App. No. 20030195154.

35 Therapeutic Methods and Administration of Antibodies

Methods provided herein comprise administering an anti-IL-4R antibody to a subject, thereby reducing an IL-4-induced biological response that plays a role in a particular condition. In particular embodiments, methods of the invention involve contacting endogenous IL-4R with an anti-IL-4R antibody, *e.g.*, in an *ex vivo* procedure.

40 Treatment encompasses alleviation or prevention of at least one symptom of a disorder, or reduction of disease severity, and the like. An antibody need not effect a complete "cure", or eradicate every symptom or manifestation of a disease, to constitute a viable therapeutic agent. As is recognized in the pertinent field, drugs employed as therapeutic agents may reduce the severity of a given disease state, but need not abolish every manifestation of the disease to be
45 regarded as useful therapeutic agents. One embodiment of the invention is directed to a method

5 comprising administering to a patient an IL-4R antagonist in an amount and for a time sufficient to induce a sustained improvement over baseline of an indicator that reflects the severity of the particular disorder.

Antibodies that inhibit the binding of both IL-4 and IL-13 to cells are discussed herein. A method for suppressing IL-4-induced and IL-13-induced activities in humans comprises
10 administering an effective amount of such an antibody. Conditions induced by IL-4 or by IL-13, or by both cytokines, thus may be treated.

As is understood in the pertinent field, antibodies are administered to a subject in a manner appropriate to the indication. Antibodies may be administered by any suitable technique, including but not limited to parenterally, topically, or by inhalation. If injected, the antibody can be
15 administered, for example, via intra-articular, intravenous, intramuscular, intralesional, intraperitoneal or subcutaneous routes, by bolus injection, or continuous infusion. Localized administration, e.g. at a site of disease or injury is contemplated, as are transdermal delivery and sustained release from implants. Delivery by inhalation includes, for example, nasal or oral inhalation, use of a nebulizer, inhalation of the antagonist in aerosol form, and the like. Other
20 alternatives include eyedrops; oral preparations including pills, syrups, lozenges or chewing gum; and topical preparations such as lotions, gels, sprays, and ointments.

Use of anti-IL-4R antibodies in *ex vivo* procedures also is contemplated. For example, a patient's blood or other bodily fluid may be contacted with an antibody that binds IL-4R *ex vivo*. The antibody may be bound to a suitable insoluble matrix or solid support material.

Advantageously, antibodies are administered in the form of a composition comprising at
25 least one anti-IL-4R antibody and one or more additional components such as a physiologically acceptable carrier, excipient or diluent. The present invention provides such compositions comprising an effective amount of an anti-IL-4R antibody, for use in the methods provided herein.

The compositions contain anti-IL-4R antibody(-ies) in, for example, any of the forms
30 described herein. The antibody may be a whole antibody or an antigen-binding fragment or engineered derivative thereof, for example.

Compositions may, for example, comprise an antibody together with a buffer, antioxidant such as ascorbic acid, low molecular weight polypeptide (such as those having fewer than 10 amino acids), protein, amino acid, carbohydrate such as glucose, sucrose or dextrans, chelating
35 agents such as EDTA, glutathione, and other stabilizers and excipients. Neutral buffered saline or saline mixed with conspecific serum albumin are examples of appropriate diluents. In accordance with appropriate industry standards, preservatives such as benzyl alcohol may also be added. The composition may be formulated as a lyophilizate using appropriate excipient solutions (e.g., sucrose) as diluents. Suitable components are nontoxic to recipients at the dosages and
40 concentrations employed. Further examples of components that may be employed in pharmaceutical formulations are presented in *Remington's Pharmaceutical Sciences*, 16th Ed., Mack Publishing Company, Easton, PA, 1980.

Kits for use by medical practitioners include an anti-IL-4R antibody and a label or other instructions for use in treating any of the conditions discussed herein. The kit preferably includes

5 a sterile preparation of one or more anti-IL-4R antibodies, which may be in the form of a composition as disclosed above, and may be in one or more vials.

Dosages and the frequency of administration may vary according to such factors as the route of administration, the particular antibody employed, the nature and severity of the disease to be treated, whether the condition is acute or chronic, and the size and general condition of the
10 subject. Appropriate dosages can be determined by procedures known in the pertinent art, e.g. in clinical trials that may involve dose escalation studies.

An antibody may be administered, for example, once or more than once, e.g., at regular intervals over a period of time. In particular embodiments, the antibody is administered over a period of at least a month or more, e.g., for one, two, or three months or even indefinitely. For
15 treating chronic conditions, long-term treatment is generally most effective. However, for treating acute conditions, administration for shorter periods, e.g. from one to six weeks, may be sufficient. In general, the antibody is administered until the patient manifests a medically relevant degree of improvement over baseline for the chosen indicator or indicators.

Particular embodiments of the present invention involve administering an antibody at a
20 dosage of from about 1 ng of antibody per kg of subject's weight per day ("1ng/kg/day") to about 10 mg/kg/day, more preferably from about 500 ng/kg/day to about 5 mg/kg/day, and most preferably from about 5 µg/kg/day to about 2 mg/kg/day, to a subject. In additional embodiments, an antibody is administered to adults one time per week, two times per week, or three or more times per week, to treat an IL-4 and/or IL-13 mediated disease, condition or disorder, e.g., a
25 medical disorder disclosed herein. If injected, the effective amount of antibody per adult dose may range from 1-20 mg/m², and preferably is about 5-12 mg/m². Alternatively, a flat dose may be administered; the amount may range from 5-100 mg/dose. One range for a flat dose is about 20-30 mg per dose. In one embodiment of the invention, a flat dose of 25 mg/dose is repeatedly administered by injection. If a route of administration other than injection is used, the dose is
30 appropriately adjusted in accordance with standard medical practices. One example of a therapeutic regimen involves injecting a dose of about 20-30 mg of anti-IL-4R antibody to one to three times per week over a period of at least three weeks, though treatment for longer periods may be necessary to induce the desired degree of improvement. For pediatric subjects (age 4-17), one exemplary suitable regimen involves the subcutaneous injection of 0.4 mg/kg, up to a
35 maximum dose of 25 mg of antibody administered two or three times per week.

Particular embodiments of the methods provided herein involve subcutaneous injection of from 0.5 mg to 10 mg, preferably from 3 to 5 mg, of an anti-IL-4R antibody, once or twice per week. Another embodiment is directed to pulmonary administration (e.g., by nebulizer) of 3 or more mg of antibody once a week.

40 Examples of therapeutic regimens provided herein comprise subcutaneous injection of an anti-IL-4R antibody once a week, at a dose of 1.5 to 3 mg, to treat asthma, pulmonary sarcoidosis, minimal change nephrosis, autoimmune uveitis, sickle cell crisis, Churg-Strauss syndrome, Sjogren's syndrome, autoimmune lymphoproliferative syndrome, pre-eclampsia, autoimmune hemolytic anemia, Barrett's esophagus, Grave's Disease, Kawasaki Disease, and cavitary
45 tuberculosis. Weekly administration of anti-IL-4R antibody is continued until a desired result is

5 achieved, *e.g.*, the subject's symptoms subside. Treatment may resume as needed, or, alternatively, maintenance doses may be administered.

10 In another embodiment, an antibody is administered to the subject in an amount and for a time sufficient to induce an improvement, preferably a sustained improvement, in at least one indicator that reflects the severity of the disorder that is being treated. Various indicators that reflect the extent of the subject's illness, disease or condition may be assessed for determining whether the amount and time of the treatment is sufficient. Such indicators include, for example, clinically recognized indicators of disease severity, symptoms, or manifestations of the disorder in question. In most instances, an improvement is considered to be sustained if the subject exhibits the improvement on at least two occasions separated by two to four weeks. The degree of improvement generally is determined by a physician, who may make this determination based on signs or symptoms, and who may also employ questionnaires that are administered to the subject, such as quality-of-life questionnaires developed for a given disease.

20 As one example, in treating benign prostate hyperplasia, an anti-IL-4R antibody is administered to the subject in an amount and for a time effective in scar regression or complete healing. Maintenance doses may be given or treatment resumed as needed.

Elevated levels of IL-4 are associated with a number of disorders, as discussed above. Subjects with a given disorder may be screened, to identify those individuals who have elevated IL-4 levels, or to identify those with an elevated TH2-type immune response, thereby identifying the subjects who may benefit most from treatment with an anti-IL-4R antibody. Thus, treatment methods provided herein optionally comprise a first step of measuring a subject's IL-4 level. An anti-IL-4R antibody may be administered to a subject in whom IL-4 levels are elevated above normal. Alternatively or additionally, a subject may be pre-screened to determine whether the subject has an elevated TH2-type immune response, prior to administration of antibody(-ies) and/or antagonist(s) against one or more TH2-type cytokines.

30 A subject's levels of IL-4 (and, optionally, of other TH2-type cytokines) may be monitored during and/or after treatment with an anti-IL-4R antibody, to detect reduction in the levels of the cytokines. For some disorders, the incidence of elevated IL-4 levels, and the balance between TH1-type and TH2-type immune responses, may vary according to such factors as the stage of the disease or the particular form of the disease. Known techniques may be employed for measuring IL-4 levels, *e.g.*, in a subject's serum, and for assessing TH2-type immune responses. Cytokine levels in blood samples may be measured by ELISA, or by a LUMINEX™ multi-plex cytokine assay (Luminex Corporation, Austin, TX) or DELFIA® (PerkinElmer LifeSciences, Wallac Oy., Turku, Finland), for example.

40 Particular embodiments of methods and compositions of the invention involve the use of an anti-IL4R antibody and one or more additional IL-4R antagonists, for example, two or more antibodies or antibody derivatives of the invention, or an antibody or antibody derivative of the invention and one or more other IL-4R antagonists. In further embodiments, anti-IL-4R antibodies are administered alone or in combination with other agents useful for treating the condition with which the patient is afflicted. Examples of such agents include both proteinaceous and non-proteinaceous drugs. When multiple therapeutics are co-administered, dosages may be adjusted

5 accordingly, as is recognized in the pertinent art. "Co-administration" and combination therapy are not limited to simultaneous administration, but include treatment regimens in which an IL-4R antibody is administered at least once during a course of treatment that involves administering at least one other therapeutic agent to the patient.

10 Examples of other agents that may be co-administered with IL-4R antibodies are other antibodies, cytokines, or cytokine receptors, which are chosen according to the particular condition to be treated. Alternatively, non-proteinaceous drugs that are useful in treating one of the particular conditions discussed above may be co-administered with an IL-4R antagonist.

15 For treating IgE-mediated conditions, an anti-IL-4R antibody may be co-administered with an IgE antagonist. One example is an anti-IgE antibody, *e.g.*, XOLAIR® (Genentech, South San Francisco, CA). Humanized anti-IgE monoclonal antibodies are described in, for example, Presta *et al.*, 1993, J. Immunol. 151:2623-32 and Adelroth *et al.*, 2000, J. Allergy Clin. Immunol. 106:253-59.

20 Anti-IL-4R antibodies may be co-administered with an IL-5 antagonist, which may be a molecule that interferes with the binding of IL-5 to an IL-5 receptor, such as an anti-IL-5R or anti-IL-5 antibody (*e.g.*, a human or humanized anti-IL-5 or anti-IL-5R monoclonal antibody), or a receptor such as a soluble human IL-5 receptor polypeptide. IL-5 has been implicated as playing a role in mediating allergic responses. Thus, administration of antagonist(s) of IL-4R and IL-5 is contemplated for treatment of allergic reactions, including but not limited to allergic asthma.

25 Further examples of agents that can be used in conjunction with IL-4R antibodies include anti-IL-4 antibodies, IL-4 muteins, IL-4 binding derivatives of IL-4R (as described in, *e.g.*, U.S. Patents 5,840,869; 5,599,905, 5,856,296, 5,767,065, 5,717,072, 6,391,581, 6,548,655, 6,472,179, and 5,844,099), IL-13 binding derivatives of IL-13, IL-4 and/or IL-13 binding chimeric derivatives of IL-4R and IL-13R, IL-13 muteins, and antagonists of CD23 (*e.g.*, anti-CD23 antibodies such as IDEC-152™ (IDEC Pharmaceuticals, San Diego, CA), Phosphodiesterase 4 (*e.g.*, ROFLUMILAST®, Byk Gulden Pharmaceuticals, Konstanz, Germany), integrins (*e.g.*, R411™, Roche, Nutley, NJ), TIMs, Gob5, STAT6, and leukotrienes.

35 For treating asthma, an IL-4R antibody may be co-administered with other anti-asthma medications, such as inhaled corticosteroids, beta agonists, leukotriene antagonists, xanthines, fluticasone, salmeterol, albuterol, non-steroidal agents such as cromolyn, and the like. IL-4R antibodies may be co-administered with other anti-allergy medications to treat allergic reactions.

40 One embodiment of the present invention is directed to co-administration of an antibody or antibody derivative of the invention and fluticasone and/or salmeterol to treat a disorder such as asthma. Compositions comprising an antibody or antibody derivative of the invention, fluticasone, and salmeterol are provided herein. ADVAIR DISKUS® (GlaxoSmithKline, Research Triangle Park, NC) comprises fluticasone propionate and salmeterol xinafoate. ADVAIR DISKUS® and the antibody or antibody derivative can be delivered by any route of administration effective for treating or preventing a particular disease, disorder, injury, or condition *e.g.*, by inhalation for treating asthma.

5 Another example of combination therapy comprises co-administration of an antibody or antibody derivative of the invention and an IL-9 antagonist to a patient who has asthma. Any suitable IL-9 antagonist may be employed, such as an IL-9 receptor (*e.g.*, a soluble form thereof), an antibody that interferes with binding of IL-9 to a cell surface receptor (*e.g.*, an antibody that binds to IL-9 or to an IL-9 receptor polypeptide), or another compound that inhibits IL-9-induced
10 biological activity. IL-9 receptors include those described in WO 93/18047 and U.S. Patents 5,789,237 and 5,962,269, which are hereby incorporated by reference herein.

In an additional embodiment of combination therapy, a method for treating ulcerative colitis comprises co-administration of an antibody or antibody derivative of the invention and at least one IL-1 antagonist. Examples of IL-1 antagonists include type I IL-1 receptor, type II IL-1
15 receptor, IL-1 receptor antagonist (IL-1Ra), antagonistic (blocking) antibodies directed against IL-1, and antagonistic antibodies directed against an IL-1 receptor. Various forms of the receptors may be employed, such as fragments, variants and fusions, for example, a soluble form of type II IL-1 receptor, *e.g.*, as described in U.S. Patent 5,350,683, hereby incorporated by reference herein.

20 One method of the present invention comprises co-administering an antibody or antibody derivative of the present invention, alone or in combination with an IL-13 antagonist(s), to a patient who has minimal change nephrosis, *e.g.*, to reduce severity of the disease.

Another method provided herein is a method for treating various allergic inflammatory conditions, comprising co-administering an antibody or antibody derivative of the invention and IL-
25 13 antagonist(s). Conditions such as asthma, allergies, and chronic lung diseases such as cystic fibrosis and chronic obstructive pulmonary disease are treated by such a method.

Any suitable IL-13 antagonist may be employed, including but not limited to IL-13 receptors (preferably soluble forms thereof), IL-13 receptor antagonists, antibodies directed against IL-13 or an IL-13R, other proteins that interfere with the binding of IL-13 to an IL-13R, and
30 compounds that inhibit IL-13-mediated signal transduction. IL-13 receptors and heterodimers comprising IL-13R polypeptides as components thereof are described above. Antibodies that are raised against IL-4R may be screened for the ability to also function as IL-13 antagonists, as discussed above.

A method for treating or preventing a condition characterized by reduced epithelial barrier function comprises co-administering an antibody or antibody derivative of the invention and one or
35 more IL-13 antagonists. Such conditions are discussed above. In one embodiment, the condition is asthma. Particular embodiments are directed to co-administering one or more antibodies or antibody derivatives of the invention and one or more IL-13 antagonists to a patient having a condition involving reduction of lung epithelial barrier function or intestinal epithelial barrier
40 function, wherein both IL-4 and IL-13 play a role in the reduced barrier function. The adverse effect of IL-13 on lung and intestinal epithelial barrier function can be confirmed using assay techniques such as those described in Example 3. See also Zund *et al.*, 1996, J. Biol. Chem. 271:7460-64.

Another method provided herein comprises co-administering an antibody or antibody
45 derivative of the invention and interferon- γ (IFN- γ) to a patient having a condition involving

5 reduction of lung epithelial barrier function. Optionally, such a method further comprises co-administering one or more IL-13 antagonists to the patient (*i.e.*, co-administering an antibody or antibody derivative of the invention, IFN- γ , and an IL-13 antagonist). In one embodiment, the patient has asthma. The antibody or antibody derivative of the invention, IFN- γ , and/or IL-13 antagonist can be administered via any method of delivery effective for treating or preventing a particular disease, disorder, condition, or injury, *e.g.*, for treating asthma, via inhalation or injection.

10 One method provided herein for treating asthma comprises administering an antibody or antibody derivative of the invention and interferon- γ to a human who has asthma. Another method for treating asthma comprises co-administering an antibody or antibody derivative of the invention, IFN- γ , and an IL-13 antagonist to a human who has asthma. In one embodiment, IFN- γ is co-administered to an asthmatic, together with an antibody that functions as an antagonist of both IL-4 and IL-13. Examples of such antibodies are described elsewhere herein.

15 A single antibody or antibody derivative of the invention may function as an IL-4 antagonist and an IL-13 antagonist, as discussed above. As an example of such an agent, some antibodies raised against IL-4R α may interfere with the binding of both IL-4 and IL-13 receptor complexes, due to the shared IL-4R α component in such multi-subunit receptor complexes (discussed above). Thus, a single antibody or antibody derivative of the invention may be employed in a method for inhibiting reduction of barrier function.

20 An antibody or antibody derivative of the invention may be co-administered with one or more leukotriene receptor antagonists to treat disorders such as allergic inflammatory diseases, *e.g.*, asthma and allergies. Examples of leukotriene receptor antagonists include but are not limited to montelukast (*e.g.*, SINGULAIR®, Merck & Co., Whitehouse, NJ), pranlukast (*e.g.*, ONON®, Ono Pharmaceuticals, Osaka, Japan), and zafirlukast (*e.g.*, ACCOLATE®, AstraZeneca, Wilmington, DE). Drugs that function as 5-lipoxygenase inhibitors may be co-administered with an IL-4R antagonist to treat asthma.

30 Methods provided herein comprise administering an antibody or antibody derivative of the invention and one or more of the following to Churg-Strauss Syndrome patients: IL-4R antagonist(s), IL-5 antagonist(s), IL-13 antagonist(s) and IgE antagonist(s). One example of such a method involves co-administering an antibody or antibody derivative of the invention and IL-5 antagonist(s) to a Churg-Strauss Syndrome patient. In another embodiment, an antibody or antibody derivative of the invention and IgE antagonist(s) are co-administered to the patient. In yet another embodiment, an antibody or antibody derivative of the invention and IL-13 antagonist(s) are co-administered to the patient.

40 The hormone relaxin may be co-administered with an antibody or antibody derivative of the invention to treat scleroderma (systemic sclerosis), idiopathic pulmonary fibrosis, or any other disorder characterized by pulmonary fibrosis, such as the conditions involving fibrosis of the lung that are discussed above. Recombinant human relaxin is preferred for use in treating humans.

A method for treating benign prostate hyperplasia comprises co-administering an antibody or antibody derivative of the invention and one or more additional anti-inflammatory agents.

5 Examples of agents that inhibit inflammation include tumor necrosis factor (TNF) antagonists and IL-17 antagonists.

Any suitable IL-17 antagonist may be employed, including but not limited to an IL-17 receptor (e.g., soluble forms thereof), IL-17 receptor antagonists, antibodies directed against IL-17 or an IL-17 receptor, other proteins that interfere with the binding of IL-17 to an IL-17 receptor, and
10 compounds that inhibit IL-17-mediated signal transduction. An IL-17 receptor, including soluble forms thereof and oligomers thereof, is described in WO 96/29408, hereby incorporated by reference. An alternative method provided herein comprises administering an IL-17 antagonist to treat a patient with benign prostate hyperplasia.

Likewise, any suitable TNF antagonist may be employed, including but not limited to a
15 TNF receptor (preferably soluble forms thereof), fusion proteins comprising a TNF receptor (or comprising the TNF-binding portion of a TNF receptor), TNF receptor antagonists, antibodies directed against TNF or a TNF receptor, other proteins that interfere with the binding of TNF to a TNF receptor, and compounds that inhibit TNF-mediated signal transduction. Further examples of TNF inhibitors are the drugs thalidomide and pentoxifylline. The TNF receptor protein known as
20 p75 or p80 TNF-R preferably is employed. A preferred TNF antagonist is a soluble human TNF receptor (sTNF-R) in dimeric form, such as dimers of sTNF-R/Fc fusion proteins. One such dimer is etanercept (ENBREL®, Immunex Corporation, Seattle, WA). p75/p80 TNF-R, including soluble fragments and other forms thereof, is described in WO 91/03553, hereby incorporated by reference herein.

25 Accordingly, in one embodiment of the present invention, an antibody or antibody derivative of the invention is co-administered with a TNF antagonist to treat any condition in which undesirable IL-4R-mediated and TNF-induced immune responses play a role, such as inflammation. One method provided herein comprises co-administering an antibody or antibody derivative of the invention and a TNF antagonist to a patient with inflammatory bowel disease,
30 Crohn's disease, or ulcerative colitis. Other embodiments are directed to a method comprising co-administering an antibody or antibody derivative of the invention and a TNF antagonist to a patient who has Kawasaki Disease, autoimmune hemolytic anemia, autoimmune uveoretinitis, autoimmune lymphoproliferative syndrome, Sjogren's syndrome, chronic fatigue syndrome, or hepatotoxicity induced by a drug such as diclofenac.

35 Another method provided herein comprises co-administering an antibody or antibody derivative of the invention and a TNF antagonist to a pregnant woman who has developed pre-eclampsia. In one embodiment, the administration of the antibody or antibody derivative of the invention TNF-antagonist continues for the duration of the pregnancy.

Suitable dosages of etanercept will vary according to the nature of the disease to be
40 treated, disease severity, the size of the patient (e.g., adult or child), and other factors, as is recognized in the pertinent field. In one embodiment of the methods provided herein, ENBREL® is administered twice a week by subcutaneous injection at a dose of from 1 to 25 mg. One embodiment of a pediatric dosage is 0.4 mg/kg. Particular methods provided herein comprise co-administration of an antibody or antibody derivative of the invention and ENBREL® to a patient has

5 autoimmune lymphoproliferative syndrome or Sjogren's syndrome, wherein ENBREL[®] is given by subcutaneous injection at a dose of from 1 to 25 mg.

For treating graft versus host disease ("GVHD"), an antibody or antibody derivative of the invention is co-administered with at least one of the following agents: a TNF antagonist, an IL-1 antagonist, steroids, or corticosteroids. In one embodiment, the TNF inhibitor is ENBREL[®]. In
10 another embodiment, the IL-1 antagonist is a soluble form of type II IL-1 receptor, *e.g.*, as described in U.S. Patent 5,350,683. In another embodiment, the GVHD is associated with (*e.g.*, develops subsequent to) bone marrow transplantation. An antibody or antibody derivative of the invention may be employed in combination with at least one of the above-listed agents, in methods
15 for suppressing an immune response directed against transplanted cells, tissue, and/or alloantigen.

A number of cytokine antagonists and other agents/drugs are disclosed herein as being useful for combination therapy (*e.g.*, co-administration with an antibody or antibody derivative of the invention) in treating particular diseases. It is to be understood that such antagonists, agents, or drugs also find use as single agents in treating those diseases. It also is to be understood that
20 disclosure of methods involving administration of an antagonist to a particular cytokine, to treat a disease, encompasses administration of one type of antagonist, and also encompasses administration of two or more different antagonists for that cytokine, unless specified otherwise.

EXAMPLE 1: Preparation of Monoclonal Antibodies

25 This example demonstrates a method of preparing monoclonal antibodies recognizing the human IL-4 receptor.

IL-4 receptor polypeptides may be employed as immunogens in generating monoclonal antibodies by conventional techniques, *e.g.*, techniques described in U.S. Patent 5,599,905, hereby incorporated by reference in its entirety. It is recognized that polypeptides in various forms
30 may be employed as immunogens, *e.g.*, full length proteins, fragments thereof, fusion proteins thereof such as Fc fusions, cells expressing the recombinant protein on the cell surface, *etc.*

To summarize an example of such a procedure, an IL-4R immunogen emulsified in complete Freund's adjuvant is injected subcutaneously into Lewis rats, in amounts ranging from 10-100 μ l. Three weeks later, the immunized animals are boosted with additional immunogen
35 emulsified in incomplete Freund's adjuvant and boosted every three weeks thereafter. Serum samples are periodically taken by retro-orbital bleeding or tail-tip excision for testing by dot-blot assay, ELISA (enzyme-linked immunosorbent assay), or inhibition of binding of ¹²⁵I-IL-4 to extracts of IL-4R-expressing cells. Following detection of an appropriate antibody titer, positive animals were given a final intravenous injection of antigen in saline. Three to four days later, the
40 animals are sacrificed, splenocytes harvested, and fused to the murine myeloma cell line AG8653. The resulting hybridoma cell lines are plated in multiple microtiter plates in a HAT selective medium (hypoxanthine, aminopterin, and thymidine) to inhibit proliferation of non-fused cells, myeloma hybrids, and spleen cell hybrids.

5 Hybridoma clones thus generated are screened for reactivity with IL-4R. Initial screening of hybridoma supernatants utilizes an antibody capture and binding of partially purified ^{125}I -IL-4 receptor. Hybridomas that are positive in this screening method are tested by a modified antibody capture to detect hybridoma cells lines that are producing blocking antibody. Hybridomas that secrete a monoclonal antibody capable of inhibiting ^{125}I -IL-4 binding to cells expressing IL-4R are
10 thus detected. Such hybridomas then are injected into the peritoneal cavities of nude mice to produce ascites containing high concentrations (>1 mg/ml) of anti-IL-4R monoclonal antibody. The resulting monoclonal antibodies may be purified by ammonium sulfate precipitation followed by gel exclusion chromatography, and/or affinity chromatography based on binding of antibody to Protein G.

15 Methods for generating human antibodies in transgenic mice have been described. See, e.g., Chen *et al.*, 1993, *Internat. Immunol.* 5: 647-56; Chen *et al.*, 1993, *EMBO J.* 12: 821-30; Choi *et al.*, 1993, *Nature Genetics* 4: 117-23; Fishwild *et al.*, 1996, *Nature Biotech.* 14: 845-51; Harding *et al.*, 1995, *Annals New York Acad. Sci.*; Lonberg *et al.*, 1994, *Nature* 368: 856-59; Lonberg, 1994, *Handbook Exper. Pharmacol.* 113: 49-101; Lonberg *et al.*, 1995, *Internat. Rev. Immunol.* 13:
20 65-93; Morrison, S, 1994, *Nature* 368: 812-13; Neuberger, 1996, *Nature Biotech.* 14: 826; Taylor *et al.*, 1992, *Nuc. Acids Res.* 20: 6287-95; Taylor *et al.*, 1994, *Internat. Immunol.* 6: 579-91; Tomizuka *et al.*, 1997, *Nature Genetics* 16: 133-43; Tomizuka *et al.*, 2000, *Proc. Nat. Acad. Sci. USA* 97: 722-27; Tuallion *et al.*, 1993, *Proc. Nat. Acad. Sci. USA* 90: 3720-24; Tuallion *et al.*, 1994, *J. Immunol.* 152: 2912-20; Russel *et al.*, 2000, *Infection and Immunity* April 2000: 1820-26;
25 Gallo *et al.*, 2000, *Eur. J. Immunol.* 30: 534-40; Davis *et al.*, 1999, *Cancer Metastasis Rev.* 18:421-25; Green, 1999, *J. Immunol. Methods* 231:11-23; Jakobovits, 1998, *Advanced Drug Delivery Rev.* 31:33-42; Green *et al.*, 1998, *J. Exp. Med.* 188: 483-95; Jakobovits, 1998, *Exp. Opin. Invest. Drugs* 7: 607-14; Tsuda *et al.*, 1997, *Genomics* 42: 413-21; Mendez *et al.*, 1997, *Nature Genetics* 15: 146-56; Jakobovits, 1996, *Weir's Handbook of Experimental Immunology*,
30 *The Integrated Immune System Vol. IV*, 194.1-194.7; Mendez *et al.*, 1995, *Genomics* 26: 294-307; Jakobovits, 1994, *Current Biol.* 4: 761-63; Arbones, 1994, *Immunity* 1: 247-60; Green *et al.*, 1994, *Nature Genetics* 7: 13-21; Jakobovits *et al.*, 1993, *Nature* 362: 255-58; Jakobovits *et al.*, 1993, *Proc. Nat. Acad. Sci. USA* 90: 2551-55.

35 **EXAMPLE 2: Assay for assessing blocking activity**

This example demonstrates an assay that can be used to identify antibodies that reduce IL-4 and/or IL-13-dependent expression of CD23. The assay is based on the ability of both IL-4 and IL-13 to enhance the expression of the activation-associated surface antigen CD23 on human B cells.

40 Antibodies raised against human IL-4R (huIL-4R) are tested either in the form of hybridoma supernatants or purified protein. Prior to addition to cultures, the antibodies are buffer exchanged against culture medium (RPMI 1640 plus 10% heat-inactivated fetal bovine serum) by centrifugation, using Centricon filter devices (Amicon) with a 10kDa cutoff.

5 Human peripheral blood B cells are purified as described previously (Morris *et al.*, 1999, J. Biol. Chem. 274:418-23). The B cells (3×10^5 /well) in culture medium are placed in 96-well round-bottomed microtiter plates and preincubated at room temperature for 30 min with test antibodies at the final concentrations indicated. Recombinant human IL-4 or IL-13 is then added to the cultures at the concentrations indicated, and cells are cultured for 20-24 hours at 37° C in a humidified atmosphere of 5% CO₂. At the end of the culture period, cells are washed once in PBS + 0.02% NaN₃ in the 96-well culture plate and are resuspended in blocking buffer (2% normal rabbit serum + 1% normal goat serum in PBS + NaN₃). Phycoerythrin (PE)-conjugated CD23 monoclonal antibody (mAb) or PE-conjugated isotype control mAb (both from Pharmingen) are then added to cells at a final dilution of 1:10. Cells are incubated for 30 minutes at 4° C, washed x3 in PBS + NaN₃ and analyzed on a FacScan (Becton Dickinson) for CD23 expression.

Cells cultured with hybridoma growth medium or isotype-matched non-blocking human anti-hIL-4R antibody are included as negative controls. An anti-huIL-4R murine mAb (MAB 230, R&D Systems, Minneapolis, MN), previously shown to block the binding and function of both hIL-4 and hIL-13, is used as a positive control for neutralization of CD23 induction by IL-4 and IL-13.

EXAMPLE 3: Assays for Measuring Loss of Barrier Function

This example provides methods of assessing the ability of an IL-4 antagonist to inhibit IL-4 induced damage to epithelial tissue and loss of epithelial barrier function.

In one aspect, the present invention provides a method of using an IL-4 antagonist to inhibit IL-4-induced damage to epithelium, including but not limited to lung epithelium or intestinal epithelium. Damage to epithelium can result in loss of barrier function. The following are non-limiting examples of techniques that may be employed in assessing the ability of an IL-4 antagonist to inhibit IL-4-induced damage to epithelium and loss of epithelial barrier function.

Cells that may be employed in preparing *in vitro* models of epithelium (epithelial barriers) are known. For example, Calu-3 human lung epithelial cells are suitable for use in barrier function studies (Ahdieh *et al.*, 2001, Am. J. Physiol. Cell Physiol. 281:C2029-38). Another suitable cell line is the human intestinal epithelial cell line designated T84. T84 cells are cultured under conditions that result in formation of a monolayer of epithelial cells on a permeable support, as described in Madara *et al.*, 1985, J. Cell Biol., 101:2124-33, Madara *et al.*, 1989, J. Clin. Invest. 83:724-27, and Youakim *et al.*, 1999, Am. J. Physiol. 276 (Gastrointest. Liver Physiol. 39):G1279-88. The thus-generated epithelial monolayer simulates the intestinal epithelial barrier.

The cultured monolayers are tested for a property (*e.g.*, resistance to passive transepithelial ion flow) that can distinguish an intact epithelium from a damaged epithelium. One such assay determines whether a particular radiolabeled compound is able to cross an epithelial monolayer. Leakage of the radiolabeled compound through the monolayer indicates that the barrier is permeable rather than intact. For example, mannitol flux analysis can be used to detect epithelial damage by assessing the movement of radiolabeled mannitol (*e.g.*, ³H mannitol) across a monolayer (see Madara and Stafford, *supra*).

5 Other examples of methods for assessing the condition of an epithelial include imaging methods (e.g., those discussed in Madara and Stafford, *supra*) and transepithelial electrical resistance measurements (e.g., those discussed in Youakim and Ahdieh, *supra*).

U.S. Patent 6,033,688 ("the '688 patent"), incorporated herein by reference in its entirety, also describes procedures that may be employed in studies of barrier permeability; see, e.g.,
10 Examples 1 and 4 of the patent. Human tracheal epithelial cells are cultured under conditions that yield a monolayer exhibiting transepithelial electrical resistance. Transepithelial resistance (indicating an intact barrier) is determined using a voltmeter. The effect of a substance or treatment on the epithelial monolayer is assessed by exposing the monolayer to the substance or treatment, and then measuring ion transport activities in Ussing chambers (column 8, lines 40-56).
15 Similar procedures can be conducted using monolayers that are generated from other types of cells, e.g., bronchial epithelial cells from a human cystic fibrosis patient (see the '688 patent, example 4, column 11).

Thus, the ability of a substance's or treatment's ability to inhibit IL-4-induced reduction in the barrier function of an epithelial layer can be assessed by exposing the epithelial layer to IL-4
20 and to the substance or treatment, assaying the condition of the epithelial layer, and comparing the condition of the epithelial layer to the condition of an epithelial layer that has been exposed to IL-4 in the absence of the substance or condition. An improvement in the condition of the epithelial layer exposed to the substance or treatment relative to an epithelial layer not exposed to the substance or treatment indicates that the substance or treatment to inhibits IL-4 induced
25 damage to epithelial tissue and loss of epithelial barrier function.

In one such assay, a monolayer of T84 cells served as an *in vitro* model of an intestinal epithelial barrier, as discussed above. IL-4 added to the basolateral side of polarized epithelial cells was found to reduce barrier function by 70% within 48-72 hours of treatment. When a soluble IL-4 receptor polypeptide consisting of the extracellular domain was added at the same
30 time as IL-4, the reduction in barrier function was prevented, and the barrier was maintained at the same level as untreated (control) cells.

The assay procedure also was conducted on a monolayer derived from lung epithelial cells, which served as an *in vitro* model of a lung epithelial barrier. IL-4 added to the basolateral side of polarized lung epithelial cells was found to reduce barrier function by 50% within 48-72
35 hours of treatment. When the IL-4 receptor polypeptide was added at the same time as IL-4, the reduction in barrier function was prevented, and the barrier was maintained at the same level as untreated (control) cells.

EXAMPLE 4: Assays for measuring binding activity

40 This example provides methods of assessing the binding activity of an anti-IL-4R antibody. Anti-huIL-4R (or a variant, derivative, or fragment thereof), is radiolabeled with ¹²⁵I using a solid phase chloramine-T analogue (IODOGEN®, Pierce, St. Louis, MO) or other, suitable radiolabeling technique, to a specific activity of approximately 3 x 10¹⁶ cpm/mmol. Loss of bioactivity is assessed by comparing binding inhibition or other biological inhibition assay with the
45 corresponding unlabeled protein. Alternatively, inhibition of radiolabeled IL-4 binding by anti-huIL-

5 4R (or a variant or fragment thereof) can be measured. Equilibrium binding assays on cells expressing IL-4R are performed in 96-well microtiter trays substantially as described in Idzerda, *et al.*, 1990, J. Exp. Med. 171:3 861-73. Briefly, serial dilutions of radiolabeled protein in binding medium (RPMI 1640, 2.5% BSA, 20 mM HEPES, 0.02% sodium azide, pH 7.2), supplemented with 0.5 mg/ml human IgG and 5% human serum), are incubated with cells (2.5×10^6 /well) for 2
10 hours at 4° C in a total volume of 150 microliters. Free and bound radiolabeled probes are separated by microfugation through a phthalate-oil separation mixture and counted in a gamma counter. Inhibition assays use radiolabeled protein at a constant concentration of 0.5 nM in the presence or absence of potential inhibitors. Nonspecific binding is determined in the presence of a 100-fold excess of unlabeled protein. Theoretical curves based on single-site competitive
15 inhibition model were fitted to the data as described in Dower *et al.*, 1984, J Immunol. 132:751. Percent inhibition is calculated according to the equation $I(\%) = [100 Ki(I)/[1 + Ka(L) + Ki(I)]]$, where I is the molar concentration of inhibitor, L is the molar concentration of radiolabeled protein, and Ki and Ka are the affinity constants of inhibitor and protein, respectively.

Equilibrium binding and competitive inhibition isotherms may also determined in 96-well
20 microtiter plates coated with IL-4R/Fc or a control Fc protein, captured through goat anti-human Fc polyclonal antibody (or other suitable anti-human Fc antibody). Briefly, plates are incubated with 5 micrograms/ml anti-human Fc in PBS at 4° C, washed twice with PBS, and then incubated with IL-4R/Fc or a control Fc protein in PBS/0.01 % Tween 20 for about 12 hours at 4° C and washed again twice with PBS. Equilibrium binding isotherms used serial dilutions of 125 I-labeled binding
25 protein in binding medium, and inhibition assays used a constant of 0.5 nM 125 I-labeled anti-huIL-4R in the presence or absence of unlabeled, potential competitive inhibitors, as described above. After 2 hours at 4° C, plates are washed twice in PBS, and specifically bound protein is released with 50 mM citrate (pH 3.0), or SDS treatment, and released 125 I-labeled anti-huIL-4R measured on a gamma counter. Data are processed as described in Dower *et al.*, *supra*.

30 Binding activity may also be assessed by surface plasmon resonance using a BIACORE® biosensor (BIAcore International AB, Uppsala, Sweden). Briefly, goat-antihuman IgG, gamma chain-specific (or other suitable gamma chain-specific antibody; GHFC) is covalently coupled to biosensor chips using a standard amine coupling procedure and reagents according to the manufacturer's instructions. Anti-huIL-4R or a control antibody is injected over the immobilized
35 GHFC, and varying amounts of IL-4R are independently passed over a GHFC coated chip (negative control) as well as an anti-huIL-4R-coated chip. Regeneration of the chip is accomplished with one 10-microliter pulse of 100 mM phosphoric acid at 10 microliters/minute. All binding is performed in HBS (10 mM HEPES, 0.15 M NaCl, 3.4 mM EDTA, 0.02 % NaN₃, 0.005 % surfactant P2O, pH 7.4).

40 **EXAMPLE 5: Anti-huIL-4R Antibodies**

This example provides the amino acid sequences, and the nucleotide sequences encoding them, of the variable domains of the heavy and light chains of antibodies that bind to huIL-4R.

5 Figure 2 provides the nucleotide sequences of the variable regions of the antibody light chains designated The light chain variable regions L1, L2, L3, L4, L5, and L6 ,and the heavy chain variable regions H1, H2, H3, H4, H5, H6, H7, H8, H9, H10, H11, H12, H13, and H14, and nucleic acids encoding them, were made. Figures 2 and 3 provide the nucleic acid and amino acid sequences, respectively, for these molecules. In Figures 2 and 3, the CDR regions of each chain are indicated in bold and underlined in L1 and H1. The FR regions also are indicated.

10 Antibodies that bind IL-4R and block binding of IL-4 and/or IL-13 thereto (e.g., scFv, Fab, F(ab')₂, diabodies, triabodies, tetrabodies, IgA, IgG1, IgG2, IgG3, IgG4, IgM, IgE, and IgD antibodies, and IgG4 antibodies having a mutation or mutations in a hinge region that alleviates a tendency to form intra-H chain disulfide bonds) can be made using combinations of the L1-L6 and
15 H1-H24 using techniques known in the art and described herein.

EXAMPLE 6: Evaluation of binding activity

This example provides binding data for certain antibodies of the invention.

Each light chain variable region L1-L6 was joined to a human kappa light chain constant region. Each heavy chain variable region H1-H14 was joined to a human IgG4 constant region.
20 Using these constructs, antibodies comprising the variable domain combinations L1H1, L2H1, L3H1, L4H1, L5H1, L1H2, L1H3, L1H4, L1H5, L1H6, L1H7, L1H8, L1H9, L1H10, L1H11, L2H4, L2H12, L2H13, L2H14, L6H1, L2H2, L2H3, L2H6, L2H7, L2H8, L2H9, L2H10, and L2H11 were made. Each of these antibodies was tested using the method of Example 2 and found to reduce
25 CD23 expression.

The foregoing examples, both working and prophetic, are non-limiting and are provided to illustrate particular embodiments of the instant invention. All references cited herein are incorporated by reference in their entireties.

CLAIMS

What is claimed is:

1. An antibody, wherein the light chain variable domain comprises a sequence of amino acids that differs from SEQ ID NO:4 only by:
 - a. at least one amino acid substitution selected from the group consisting of S28T, S30N, S30G, S31N, S32D, S32N, A52T, S54Y, T57P, T57S, G93D, S94H, S94R, P96A, P97G, and T99M,
 - and
 - b. optionally, one or more amino acid substitutions selected from the group consisting of E1D, L4M, S7T, G9A, K40R, F50Y, S68F, S77T, V86I, K105R, V106L, and E107D, and wherein the heavy chain variable domain comprises a sequence of amino acids that differs from SEQ ID NO:16 only by:
 - c. at least one amino acid substitution selected from the group consisting of N58S, Y101W, F102Y, D103T, D103N, D103P, Y104H, Y104N, Y104W, and Y104R,
 - and
 - d. optionally, one or more amino acid substitutions selected from the group consisting of Q6E, H13Q, G24A, R86S, and M90T,wherein said antibody binds to IL-4 receptor alpha.
2. The antibody of Claim 1, wherein:
 - a. the light chain CDR1 comprises a sequence selected from the group consisting of residues 24-35 of SEQ ID NO:6, residues 24-35 of SEQ ID NO:8, residues 24-35 of SEQ ID NO:10, residues 24-35 of SEQ ID NO:12; and residues 24-35 of SEQ ID NO:14,
 - b. the light chain CDR2 comprises a sequence selected from the group consisting of residues 51-57 of SEQ ID NO:4, residues 51-57 of SEQ ID NO:6, residues 51-57 of SEQ ID NO:10, and residues 51-57 of SEQ ID NO:12,
 - c. the light chain CDR3 comprises a sequence selected from the group consisting of residues 90-99 of SEQ ID NO:4, residues 90-99 of SEQ ID NO:6, residues 90-99 of SEQ ID NO:8, and residues 90-99 of SEQ ID NO:14,
 - d. the heavy chain CDR1 comprises the sequence residues 31-35 of SEQ ID NO:16,
 - e. the heavy chain CDR2 comprises a sequence selected from the group consisting of residues 50-65 of SEQ ID NO:16, and residues 50-65 of SEQ ID NO:18, and
 - f. the heavy chain CDR3 comprises a sequence selected from the group consisting of residues 90-99 of SEQ ID NO:4, residues 98-104 of SEQ ID NO:18, residues 98-104 of SEQ ID NO:20, residues 98-104 of SEQ ID NO:22, residues 98-104 of SEQ ID NO:24, residues 98-104 of SEQ ID NO:26, residues 98-104 of SEQ ID NO:30, and residues 98-104 of SEQ ID NO:34.

3. The antibody of Claim 2 wherein:
 - a. the light chain FR1 comprises a sequence selected from the group consisting of residues 1-23 of SEQ ID NO:4, residues 1-23 of SEQ ID NO:10, residues 1-23 of SEQ ID NO:12, and residues 1-23 of SEQ ID NO:14,
 - b. the light chain FR2 comprises a sequence selected from the group consisting of residues 36-50 of SEQ ID NO:4, and residues 36-50 of SEQ ID NO:14,
 - c. the light chain FR3 comprises a sequence selected from the group consisting of residues 58-89 of SEQ ID NO:4, residues 58-89 of SEQ ID NO:10, and residues 58-89 of SEQ ID NO:12,
 - d. the light chain FR4 comprises a sequence selected from the group consisting of residues 100-109 of SEQ ID NO:4, residues 100-109 of SEQ ID NO:8, and residues 100-109 of SEQ ID NO:12,
 - e. the heavy chain FR1 comprises a sequence selected from the group consisting of residues 1-30 of SEQ ID NO:16, and residues 1-30 of SEQ ID NO:42,
 - f. the heavy chain FR2 comprises the sequence of residues 36-49 of SEQ ID NO:16,
 - g. the heavy chain FR3 comprises a sequence selected from the group consisting of residues 66-97 of SEQ ID NO:16, residues 66-97 of SEQ ID NO:18, and residues 66-97 of SEQ ID NO:42, and
 - h. the heavy chain FR4 comprises the sequence of residues 105-115 of SEQ ID NO:16.
4. An antibody comprising:
 - a. a light chain variable domain comprising a sequence that is at least 80% identical to a sequence selected from the group consisting of SEQ ID NO:4, 6, 8, 10, 12, and 14, with the proviso that said light chain variable domain does not comprise the sequence of SEQ ID NO:4, and
 - b. a heavy chain variable domain comprising a sequence that is at least 80% identical to a sequence selected from the group consisting of SEQ ID NO:16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, and 62, with the proviso that said heavy chain variable domain does not comprise the sequence of SEQ ID NO:16,wherein said antibody binds to IL-4 receptor alpha.
5. The antibody of Claim 4, wherein:
 - a. said light chain variable domain comprises a sequence that is at least 85% identical to a sequence selected from the group consisting of SEQ ID NO:4, 6, 8, 10, 12, and 14, and
 - b. said heavy chain variable domain comprises a sequence that is at least 85% identical to a sequence selected from the group consisting of SEQ ID NO:16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, and 62.
6. The antibody of Claim 5, wherein:

- a. said light chain variable domain comprises a sequence that is at least 90% identical to a sequence selected from the group consisting of SEQ ID NO:4, 6, 8, 10, 12, and 14, and
 - b. said heavy chain variable domain comprises a sequence that is at least 90% identical to a sequence selected from the group consisting of SEQ ID NO:16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, and 62.
- 7. The antibody of Claim 6, wherein:
 - a. said light chain variable domain comprises a sequence that is at least 95% identical to a sequence selected from the group consisting of SEQ ID NO:4, 6, 8, 10, 12, and 14, and
 - b. said heavy chain variable domain comprises a sequence that is at least 95% identical to a sequence selected from the group consisting of SEQ ID NO:16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, and 62.
- 8. The antibody of Claim 7, wherein:
 - a. said light chain variable domain comprises a sequence that is selected from the group consisting of SEQ ID NO:4, 6, 8, 10, 12, and 14, and
 - b. said heavy chain variable domain comprises a sequence that is selected from the group consisting of SEQ ID NO:16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, and 62.
- 9. An antibody comprising:
 - a. a light chain variable domain comprising a sequence of at least 15 contiguous amino acids of a sequence selected from the group consisting of SEQ ID NO:4, 6, 8, 10, 12, and 14, and/or
 - b. a heavy chain variable domain comprising a sequence of at least 15 contiguous amino acids of a sequence selected from the group consisting of SEQ ID NO:16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, and 62,with the proviso that said antibody does not comprise both the light chain variable domain sequence of SEQ ID NO:4 and the heavy chain variable domain sequence of SEQ ID NO:16, and wherein said antibody binds to IL-4 receptor alpha.
- 10. The antibody of Claim 9 wherein
 - a. said light chain variable domain comprises a sequence of at least 20 contiguous amino acids of a sequence selected from the group consisting of: SEQ ID NO:4, 6, 8, 10, 12, and 14, and/or
 - b. said heavy chain variable domain comprises a sequence of at least 20 contiguous amino acids of a sequence selected from the group consisting of SEQ ID NO:16, 18,

20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, and 62.

11. The antibody of Claim 10 wherein
 - a. said light chain variable domain comprises a sequence of at least 25 contiguous amino acids of a sequence selected from the group consisting of SEQ ID NO:4, 6, 8, 10, 12, and 14, and/or
 - b. said heavy chain variable domain comprises a sequence of at least 25 contiguous amino acids of a sequence selected from the group consisting of SEQ ID NO:16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, and 62.
12. The antibody of Claim 11 wherein
 - a. said light chain variable domain comprises a sequence of at least 35 contiguous amino acids of a sequence selected from the group consisting of SEQ ID NO:4, 6, 8, 10, 12, and 14, and/or
 - b. said heavy chain variable domain comprises a sequence of at least 35 contiguous amino acids of a sequence selected from the group consisting of SEQ ID NO:16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, and 62.
13. The antibody of Claim 12 wherein
 - a. said light chain variable domain comprises a sequence of at least 50 contiguous amino acids of a sequence selected from the group consisting of SEQ ID NO:4, 6, 8, 10, 12, and 14, and/or
 - b. said heavy chain variable domain comprises a sequence of at least 50 contiguous amino acids of a sequence selected from the group consisting of SEQ ID NO:16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, and 62.
14. The antibody of Claim 13 wherein
 - a. said light chain variable domain comprises a sequence of at least 75 contiguous amino acids of a sequence selected from the group consisting of SEQ ID NO:4, 6, 8, 10, 12, and 14, and/or
 - b. said heavy chain variable domain comprises a sequence of at least 75 contiguous amino acids of a sequence selected from the group consisting of SEQ ID NO:16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, and 62.
15. The antibody of Claim 14 wherein

- a. said light chain variable domain comprises a sequence of at least 100 contiguous amino acids of a sequence selected from the group consisting of SEQ ID NO:4, 6, 8, 10, 12, and 14, and/or
 - b. said heavy chain variable domain comprises a sequence of at least 100 contiguous amino acids of a sequence selected from the group consisting of SEQ ID NO:16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, and 62.
- 16. An antibody comprising:
 - a. a light chain variable domain comprising an amino acid sequence that is encoded by a nucleotide sequence that is at least 80% identical to a nucleotide sequence selected from the group consisting of SEQ ID NO:3, 5, 7, 9, 11, and 13, and/or
 - b. a heavy chain variable domain comprising an amino acid sequence that is encoded by a nucleotide sequence that is at least 80% identical to a nucleotide sequence selected from the group consisting of SEQ ID NO:15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59, and 61.with the proviso that said antibody does not comprise both the light chain variable domain sequence of SEQ ID NO:4 and the heavy chain variable domain sequence of SEQ ID NO:16, and wherein said antibody binds to IL-4 receptor alpha.
- 17. The antibody of Claim 16, wherein:
 - a. said light chain variable domain comprises an amino acid sequence that is encoded by a nucleotide sequence that is at least 85% identical to a nucleotide sequence selected from the group consisting of SEQ ID NO:3, 5, 7, 9, 11, and 13, and/or
 - b. said heavy chain variable domain comprises an amino acid sequence that is encoded by a nucleotide sequence that is at least 85% identical to a nucleotide sequence selected from the group consisting of SEQ ID NO:15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59, and 61.
- 18. The antibody of Claim 17, wherein:
 - a. said light chain variable domain comprises an amino acid sequence that is encoded by a nucleotide sequence that is at least 90% identical to a nucleotide sequence selected from the group consisting of SEQ ID NO:3, 5, 7, 9, 11, and 13, and/or
 - b. said heavy chain variable domain comprises an amino acid sequence that is encoded by a nucleotide sequence that is at least 90% identical to a nucleotide sequence selected from the group consisting of SEQ ID NO:15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59, and 61.
- 19. The antibody of Claim 18, wherein:

- a. said light chain variable domain comprises an amino acid sequence that is encoded by a nucleotide sequence that is at least 95% identical to a nucleotide sequence selected from the group consisting SEQ ID NO:3, 5, 7, 9, 11, and 13, and/or
 - b. said heavy chain variable domain comprises an amino acid sequence that is encoded by a nucleotide sequence that is at least 95% identical to a nucleotide sequence selected from the group consisting of SEQ ID NO:15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59, and 61.
- 20. The antibody of Claim 19, wherein:
 - a. said light chain variable domain comprises an amino acid sequence that is encoded by a nucleotide sequence that is at least 98% identical to a nucleotide sequence selected from the group consisting of SEQ ID NO:3, 5, 7, 9, 11, and 13, and/or
 - b. said heavy chain variable domain comprises an amino acid sequence that is encoded by a nucleotide sequence that is at least 98% identical to a nucleotide sequence selected from the group consisting of SEQ ID NO:15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59, and 61.
- 21. The antibody of Claim 20, wherein:
 - a. said light chain variable domain comprises an amino acid sequence that is encoded by a nucleotide sequence selected from the group consisting of SEQ ID NO:3, 5, 7, 9, 11, and 13, and/or
 - b. said heavy chain variable domain comprises an amino acid sequence that is encoded by a nucleotide sequence selected from the group consisting of SEQ ID NO:15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59, and 61.
- 22. An antibody comprising:
 - a. a light chain variable domain comprising an amino acid sequence that is encoded by a nucleotide sequence that hybridizes under moderately stringent conditions to the complement of a nucleotide sequence selected from the group consisting of SEQ ID NO:3, 5, 7, 9, 11, and 13, and/or
 - b. a heavy chain variable domain comprising an amino acid sequence that is encoded by a nucleotide sequence that hybridizes under moderately stringent conditions to the complement of a nucleotide sequence selected from the group consisting of SEQ ID NO:15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59, and 61,with the proviso that said antibody does not comprise both the light chain variable domain sequence of SEQ ID NO:4 and the heavy chain variable domain sequence of SEQ ID NO:16, and wherein said antibody binds to IL-4 receptor alpha.
- 23. The antibody of Claim 22, wherein:

- a. said light chain variable domain comprises an amino acid sequence that is encoded by a nucleotide sequence that hybridizes under stringent conditions to the complement of a nucleotide sequence selected from the group consisting of SEQ ID NO:3, 5, 7, 9, 11, and 13, and/or
 - b. said heavy chain variable domain comprises an amino acid sequence that is encoded by a nucleotide sequence that hybridizes under stringent conditions to the complement of a nucleotide sequence selected from the group consisting of SEQ ID NO:15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59, and 61.
24. The antibody of Claim 1, wherein said antibody is selected from the group consisting of L2H1, L3H1, L4H1, L5H1, L1H2, L1H3, L1H4, L1H5, L1H6, L1H7, L1H8, L1H9, L1H10, L1H11, L2H4, L2H12, L2H13, L2H14, L6H1, L2H2, L2H3, L2H6, L2H7, L2H8, L2H9, L2H10, and L2H11.
25. The antibody of Claim 1, 4, 9, 16, or 22 wherein said antibody is a human, humanized, or chimeric antibody.
26. The antibody of Claim 1, 4, 9, 16, or 22 wherein said antibody is a monoclonal antibody.
27. The antibody of Claim 1, 4, 9, 16, or 22 wherein said antibody is selected from the group consisting of an IgD, IgE, IgM, IgG1, IgG2, IgG3, IgG4, and IgG4 having at least one mutation in a hinge region that alleviates a tendency to form intra-H chain disulfide bond antibody.
28. An isolated polypeptide comprising an IL-4 receptor binding portion of the antibody of Claim 1, 4, 9, 16, or 22.
29. The isolated polypeptide of Claim 28, wherein said polypeptide comprises a Fab, F(ab')₂, scFv, diabody, triabody, or tetrabody.
30. An isolated nucleic acid comprising a nucleotide sequence encoding the light chain of the antibody of Claim 1, 4, 9, 16, or 22.
31. The isolated nucleic acid of Claim 30, wherein said nucleic acid comprises a nucleotide sequence selected from the group consisting of SEQ ID NO:5, 7, 9, 11, and 13.
32. A vector comprising the isolated nucleic acid of Claim 30.
33. The vector of Claim 32 wherein said vector is an expression vector.

34. An isolated nucleic acid comprising a nucleotide sequence encoding the heavy chain of the antibody of Claim 1, 4, 9, 16, or 22.
35. The isolated nucleic acid of Claim 34, wherein said nucleic acid comprises a nucleotide sequence selected from the group consisting of SEQ ID NO:15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59, and 61.
36. A vector comprising the isolated nucleic acid of Claim 34.
37. The vector of Claim 36 wherein said vector is an expression vector.
38. An isolated nucleic acid comprising a nucleotide sequence encoding the isolated polypeptide of Claim 28.
39. The isolated nucleic acid of Claim 38, wherein said nucleic acid comprises a nucleotide sequence selected from the group consisting of SEQ ID NO:5, 7, 9, 11, 13, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59, and 61.
40. A vector comprising the isolated nucleic acid of Claim 38.
41. The vector of Claim 40 wherein said vector is an expression vector.
42. An isolated cell comprising the isolated nucleic acid of Claim 30.
43. The isolated cell of Claim 42 wherein said cell is a hybridoma.
44. The isolated cell of Claim 42 wherein said cell is a transgenic cell.
45. An isolated cell comprising the isolated nucleic acid of Claim 34.
46. The isolated cell of Claim 43 wherein said cell is a hybridoma.
47. The isolated cell of Claim 43 wherein said cell is a transgenic cell.
48. An isolated cell comprising the isolated nucleic acid of Claim 38.
49. The isolated cell of Claim 48 wherein said cell is a transgenic cell.
50. An isolated cell comprising the antibody of Claim 1, 4, 9, 16, or 22.
51. The isolated cell of Claim 50 wherein said cell is a hybridoma.

52. The isolated cell of Claim 50 wherein said cell is a transgenic cell.
53. A method of making the antibody of Claim 1, 4, 9, 16, or 22 comprising incubating a cell comprising a nucleic acid encoding the light chain of said antibody and a nucleic acid encoding the heavy chain of said antibody under conditions that allow said cell to express said light chain and said heavy chain and that allow said light chain and said heavy chain to assemble into said antibody, and isolating said antibody from said cell.
54. The method of Claim 53, wherein said cell is a hybridoma.
55. The method of Claim 53, wherein said cell is a transgenic cell.
56. A method of inhibiting an IL-4 receptor comprising contacting a cell expressing IL-4 receptor alpha with the antibody of Claim 1, 4, 9, 16, or 22 under conditions that allow said antibody to bind to said IL-4 receptor alpha.
57. The method of Claim 56 wherein said cell is a human cell.
58. The method of Claim 57 wherein said human cell is in a human.
59. A method of inhibiting an IL-4 receptor comprising contacting a cell expressing IL-4 receptor alpha with the polypeptide of Claim 28 under conditions that allow said polypeptide to bind to said IL-4 receptor alpha.
60. The method of Claim 59 wherein said cell is a human cell.
61. The method of Claim 60 wherein said human cell is in a human.
62. A method of treating a condition in a subject comprising administering to said subject an amount of the antibody of Claim 1, 4, 9, 16, or 22 effective for treating said condition.
63. The method of Claim 62 wherein said condition is an inflammatory or cancerous condition.
64. The method of Claim 63 wherein said inflammatory or cancerous condition is an immunological condition.
65. The method of Claim 62 wherein said condition is asthma, septic arthritis, dermatitis herpetiformis, chronic idiopathic urticaria, ulcerative colitis, scleroderma, hypertrophic scarring, Whipple's Disease, benign prostate hyperplasia, a lung disorder in which IL-4 receptor plays a role, condition in which IL-4 receptor-mediated epithelial barrier disruption

plays a role, a disorder of the digestive system in which IL-4 receptor plays a role, an allergic reaction to a medication, Kawasaki disease, sickle cell disease, Churg-Strauss syndrome, Grave's disease, pre-eclampsia, Sjogren's syndrome, autoimmune lymphoproliferative syndrome, autoimmune hemolytic anemia, Barrett's esophagus, autoimmune uveitis, tuberculosis, cystic fibrosis, allergic bronchopulmonary mycosis, chronic obstructive pulmonary disease, bleomycin-induced pneumopathy and fibrosis, radiation-induced pulmonary fibrosis, pulmonary alveolar proteinosis, adult respiratory distress syndrome, sarcoidosis, hyper IgE syndrome, idiopathic hypereosinophil syndrome, an autoimmune blistering disease, pemphigus vulgaris, bullous pemphigoid, myasthenia gravis, chronic fatigue syndrome, or nephrosis.

66. A method of treating a condition in a subject comprising administering to said subject an amount of the polypeptide of Claim 28 effective for treating said condition.
67. The method of Claim 66 wherein said condition is an inflammatory or cancerous condition.
68. The method of Claim 67 wherein said inflammatory or cancerous condition is an immunological condition.
69. The method of Claim 66 wherein said condition is asthma, septic arthritis, dermatitis herpetiformis, chronic idiopathic urticaria, ulcerative colitis, scleroderma, hypertrophic scarring, Whipple's Disease, benign prostate hyperplasia, a lung disorder in which IL-4 receptor plays a role, condition in which IL-4 receptor-mediated epithelial barrier disruption plays a role, a disorder of the digestive system in which IL-4 receptor plays a role, an allergic reaction to a medication, Kawasaki disease, sickle cell disease, Churg-Strauss syndrome, Grave's disease, pre-eclampsia, Sjogren's syndrome, autoimmune lymphoproliferative syndrome, autoimmune hemolytic anemia, Barrett's esophagus, autoimmune uveitis, tuberculosis, cystic fibrosis, allergic bronchopulmonary mycosis, chronic obstructive pulmonary disease, bleomycin-induced pneumopathy and fibrosis, radiation-induced pulmonary fibrosis, pulmonary alveolar proteinosis, adult respiratory distress syndrome, sarcoidosis, hyper IgE syndrome, idiopathic hypereosinophil syndrome, an autoimmune blistering disease, pemphigus vulgaris, bullous pemphigoid, myasthenia gravis, chronic fatigue syndrome, or nephrosis.
70. A pharmaceutical composition comprising an antibody of Claim 1, 4, 9, 16, or 22 and an excipient, diluent, or buffer.
71. A pharmaceutical composition comprising a polypeptide of Claim 28 and an excipient, diluent, or buffer.

ABSTRACT OF THE DISCLOSURE

The present invention relates to antibodies that bind to the IL-4 receptor, fragments, muteins, and derivatives of such antibodies, nucleic acids encoding such antibodies, fragments, muteins and derivatives, and methods of making and using such antibodies, fragments, muteins, derivatives and nucleic acids. Methods for treating medical conditions induced by interleukin-4 involve administering an IL-4 receptor binding antibody, or an IL-4 receptor binding fragment, mutein, or derivative of an IL-4 receptor binding antibody, to a patient afflicted with such a condition. Particular antibodies provided herein include human monoclonal antibodies. Certain of the antibodies inhibit both IL-4-induced and IL-13-induced biological activities.

Title: ANTIBODIES THAT BIND INTERLEUKIN-4
RECEPTOR

Filed: November 7, 2003

Inventors: Hongxing Zhou et al.

Docket No.: 3492

FIGURE 1A

ATG	GGG	TGG	CTT	TGC	TCT	GGG	CTC	CTG	TTC	CCT	GTG	AGC	TGC	CTG	-31
Met	Gly	Trp	Leu	Cys	Ser	Gly	Leu	Leu	Phe	Pro	Val	Ser	Cys	Leu	-11
GTC	CTG	CTG	CAG	GTG	GCA	AGC	TCT	GGG	AAC	ATG	AAG	GTC	TTG	CAG	15
Val	Leu	Leu	Gln	Val	Ala	Ser	Ser	Gly	Asn	Met	Lys	Val	Leu	Gln	5
GAG	CCC	ACC	TGC	GTC	TCC	GAC	TAC	ATG	AGC	ATC	TCT	ACT	TGC	GAG	60
Glu	Pro	Thr	Cys	Val	Ser	Asp	Tyr	Met	Ser	Ile	Ser	Thr	Cys	Glu	20
TGG	AAG	ATG	AAT	GGT	CCC	ACC	AAT	TGC	AGC	ACC	GAG	CTC	CGC	CTG	105
Trp	Lys	Met	Asn	Gly	Pro	Thr	Asn	Cys	Ser	Thr	Glu	Leu	Arg	Leu	35
TTG	TAC	CAG	CTG	GTT	TTT	CTG	CTC	TCC	GAA	GCC	CAC	ACG	TGT	ATC	150
Leu	Tyr	Gln	Leu	Val	Phe	Leu	Leu	Ser	Glu	Ala	His	Thr	Cys	Ile	50
CCT	GAG	AAC	AAC	GGA	GGC	GCG	GGG	TGC	GTG	TGC	CAC	CTG	CTC	ATG	195
Pro	Glu	Asn	Asn	Gly	Gly	Ala	Gly	Cys	Val	Cys	His	Leu	Leu	Met	65
GAT	GAC	GTG	GTC	AGT	GCG	GAT	AAC	TAT	ACA	CTG	GAC	CTG	TGG	GCT	240
Asp	Asp	Val	Val	Ser	Ala	Asp	Asn	Tyr	Thr	Leu	Asp	Leu	Trp	Ala	80
GGG	CAG	CAG	CTG	CTG	TGG	AAG	GGC	TCC	TTC	AAG	CCC	AGC	GAG	CAT	285
Gly	Gln	Gln	Leu	Leu	Trp	Lys	Gly	Ser	Phe	Lys	Pro	Ser	Glu	His	95
GTG	AAA	CCC	AGG	GCC	CCA	GGA	AAC	CTG	ACA	GTT	CAC	ACC	AAT	GTC	330
Val	Lys	Pro	Arg	Ala	Pro	Gly	Asn	Leu	Thr	Val	His	Thr	Asn	Val	110
TCC	GAC	ACT	CTG	CTG	CTG	ACC	TGG	AGC	AAC	CCG	TAT	CCC	CCT	GAC	375
Ser	Asp	Thr	Leu	Leu	Leu	Thr	Trp	Ser	Asn	Pro	Tyr	Pro	Pro	Asp	125
AAT	TAC	CTG	TAT	AAT	CAT	CTC	ACC	TAT	GCA	GTC	AAC	ATT	TGG	AGT	420
Asn	Tyr	Leu	Tyr	Asn	His	Leu	Thr	Tyr	Ala	Val	Asn	Ile	Trp	Ser	140
GAA	AAC	GAC	CCG	GCA	GAT	TTC	AGA	ATC	TAT	AAC	GTG	ACC	TAC	CTA	465
Glu	Asn	Asp	Pro	Ala	Asp	Phe	Arg	Ile	Tyr	Asn	Val	Thr	Tyr	Leu	155
GAA	CCC	TCC	CTC	CGC	ATC	GCA	GCC	AGC	ACC	CTG	AAG	TCT	GGG	ATT	510
Glu	Pro	Ser	Leu	Arg	Ile	Ala	Ala	Ser	Thr	Leu	Lys	Ser	Gly	Ile	170
TCC	TAC	AGG	GCA	CGG	GTG	AGG	GCC	TGG	GCT	CAG	TGC	TAT	AAC	ACC	555
Ser	Tyr	Arg	Ala	Arg	Val	Arg	Ala	Trp	Ala	Gln	Cys	Tyr	Asn	Thr	185
ACC	TGG	AGT	GAG	TGG	AGC	CCC	AGC	ACC	AAG	TGG	CAC	AAC	TCC	TAC	600
Thr	Trp	Ser	Glu	Trp	Ser	Pro	Ser	Thr	Lys	Trp	His	Asn	Ser	Tyr	200
AGG	GAG	CCC	TTC	GAG	CAG	CAC	CTC	CTG	CTG	GGC	GTC	AGC	GTT	TCC	645
Arg	Glu	Pro	Phe	Glu	Gln	His	Leu	Leu	Leu	Gly	Val	Ser	Val	Ser	215
TGC	ATT	GTC	ATC	CTG	GCC	GTC	TGC	CTG	TTG	TGC	TAT	GTC	AGC	ATC	690
Cys	Ile	Val	Ile	Leu	Ala	Val	Cys	Leu	Leu	Cys	Tyr	Val	Ser	Ile	230
ACC	AAG	ATT	AAG	AAA	GAA	TGG	TGG	GAT	CAG	ATT	CCC	AAC	CCA	GCC	735
Thr	Lys	Ile	Lys	Lys	Glu	Trp	Trp	Asp	Gln	Ile	Pro	Asn	Pro	Ala	245

FIGURE 1B

CGC	AGC	CGC	CTC	GTG	GCT	ATA	ATA	ATC	CAG	GAT	GCT	CAG	GGG	TCA	780
Arg	Ser	Arg	Leu	Val	Ala	Ile	Ile	Ile	Gln	Asp	Ala	Gln	Gly	Ser	260
CAG	TGG	GAG	AAG	CGG	TCC	CGA	GGC	CAG	GAA	CCA	GCC	AAG	TGC	CCA	825
Gln	Trp	Glu	Lys	Arg	Ser	Arg	Gly	Gln	Glu	Pro	Ala	Lys	Cys	Pro	275
CAC	TGG	AAG	AAT	TGT	CTT	ACC	AAG	CTC	TTG	CCC	TGT	TTT	CTG	GAG	870
His	Trp	Lys	Asn	Cys	Leu	Thr	Lys	Leu	Leu	Pro	Cys	Phe	Leu	Glu	290
CAC	AAC	ATG	AAA	AGG	GAT	GAA	GAT	CCT	CAC	AAG	GCT	GCC	AAA	GAG	915
His	Asn	Met	Lys	Arg	Asp	Glu	Asp	Pro	His	Lys	Ala	Ala	Lys	Glu	305
ATG	CCT	TTC	CAG	GGC	TCT	GGA	AAA	TCA	GCA	TGG	TGC	CCA	GTG	GAG	960
Met	Pro	Phe	Gln	Gly	Ser	Gly	Lys	Ser	Ala	Trp	Cys	Pro	Val	Glu	320
ATC	AGC	AAG	ACA	GTC	CTC	TGG	CCA	GAG	AGC	ATC	AGC	GTG	GTG	CGA	1005
Ile	Ser	Lys	Thr	Val	Leu	Trp	Pro	Glu	Ser	Ile	Ser	Val	Val	Arg	335
TGT	GTG	GAG	TTG	TTT	GAG	GCC	CCG	GTG	GAG	TGT	GAG	GAG	GAG	GAG	1050
Cys	Val	Glu	Leu	Phe	Glu	Ala	Pro	Val	Glu	Cys	Glu	Glu	Glu	Glu	350
GAG	GTA	GAG	GAA	GAA	AAA	GGG	AGC	TTC	TGT	GCA	TCG	CCT	GAG	AGC	1095
Glu	Val	Glu	Glu	Glu	Lys	Gly	Ser	Phe	Cys	Ala	Ser	Pro	Glu	Ser	365
AGC	AGG	GAT	GAC	TTC	CAG	GAG	GGA	AGG	GAG	GGC	ATT	GTG	GCC	CGG	1140
Ser	Arg	Asp	Asp	Phe	Gln	Glu	Gly	Arg	Glu	Gly	Ile	Val	Ala	Arg	380
CTA	ACA	GAG	AGC	CTG	TTC	CTG	GAC	CTG	CTC	GGA	GAG	GAG	AAT	GGG	1185
Leu	Thr	Glu	Ser	Leu	Phe	Leu	Asp	Leu	Leu	Gly	Glu	Glu	Asn	Gly	395
GGC	TTT	TGC	CAG	CAG	GAC	ATG	GGG	GAG	TCA	TGC	CTT	CTT	CCA	CCT	1230
Gly	Phe	Cys	Gln	Gln	Asp	Met	Gly	Glu	Ser	Cys	Leu	Leu	Pro	Pro	410
TCG	GGA	AGT	ACG	AGT	GCT	CAC	ATG	CCC	TGG	GAT	GAG	TTC	CCA	AGT	1275
Ser	Gly	Ser	Thr	Ser	Ala	His	Met	Pro	Trp	Asp	Glu	Phe	Pro	Ser	425
GCA	GGG	CCC	AAG	GAG	GCA	CCT	CCC	TGG	GGC	AAG	GAG	CAG	CCT	CTC	1320
Ala	Gly	Pro	Lys	Glu	Ala	Pro	Pro	Trp	Gly	Lys	Glu	Gln	Pro	Leu	440
CAC	CTG	GAG	CCA	AGT	CCT	CCT	GCC	AGC	CCG	ACC	CAG	AGT	CCA	GAC	1365
His	Leu	Glu	Pro	Ser	Pro	Pro	Ala	Ser	Pro	Thr	Gln	Ser	Pro	Asp	455
AAC	CTG	ACT	TGC	ACA	GAG	ACG	CCC	CTC	GTC	ATC	GCA	GGC	AAC	CCT	1410
Asn	Leu	Thr	Cys	Thr	Glu	Thr	Pro	Leu	Val	Ile	Ala	Gly	Asn	Pro	470
GCT	TAC	CGC	AGC	TTC	AGC	AAC	TCC	CTG	AGC	CAG	TCA	CCG	TGT	CCC	1455
Ala	Tyr	Arg	Ser	Phe	Ser	Asn	Ser	Leu	Ser	Gln	Ser	Pro	Cys	Pro	485
AGA	GAG	CTG	GGT	CCA	GAC	CCA	CTG	CTG	GCC	AGA	CAC	CTG	GAG	GAA	1500
Arg	Glu	Leu	Gly	Pro	Asp	Pro	Leu	Leu	Ala	Arg	His	Leu	Glu	Glu	500
GTA	GAA	CCC	GAG	ATG	CCC	TGT	GTC	CCC	CAG	CTC	TCT	GAG	CCA	ACC	1545
Val	Glu	Pro	Glu	Met	Pro	Cys	Val	Pro	Gln	Leu	Ser	Glu	Pro	Thr	515

FIGURE 1C

ACT	GTG	CCC	CAA	CCT	GAG	CCA	GAA	ACC	TGG	GAG	CAG	ATC	CTC	CGC	1590
Thr	Val	Pro	Gln	Pro	Glu	Pro	Glu	Thr	Trp	Glu	Gln	Ile	Leu	Arg	530
CGA	AAT	GTC	CTC	CAG	CAT	GGG	GCA	GCT	GCA	GCC	CCC	GTC	TCG	GCC	1635
Arg	Asn	Val	Leu	Gln	His	Gly	Ala	Ala	Ala	Ala	Pro	Val	Ser	Ala	545
CCC	ACC	AGT	GGC	TAT	CAG	GAG	TTT	GTA	CAT	GCG	GTG	GAG	CAG	GGT	1680
Pro	Thr	Ser	Gly	Tyr	Gln	Glu	Phe	Val	His	Ala	Val	Glu	Gln	Gly	560
GGC	ACC	CAG	GCC	AGT	GCG	GTG	GTG	GGC	TTG	GGT	CCC	CCA	GGA	GAG	1725
Gly	Thr	Gln	Ala	Ser	Ala	Val	Val	Gly	Leu	Gly	Pro	Pro	Gly	Glu	575
GCT	GGT	TAC	AAG	GCC	TTC	TCA	AGC	CTG	CTT	GCC	AGC	AGT	GCT	GTG	1770
Ala	Gly	Tyr	Lys	Ala	Phe	Ser	Ser	Leu	Leu	Ala	Ser	Ser	Ala	Val	590
TCC	CCA	GAG	AAA	TGT	GGG	TTT	GGG	GCT	AGC	AGT	GGG	GAA	GAG	GGG	1815
Ser	Pro	Glu	Lys	Cys	Gly	Phe	Gly	Ala	Ser	Ser	Gly	Glu	Glu	Gly	605
TAT	AAG	CCT	TTC	CAA	GAC	CTC	ATT	CCT	GGC	TGC	CCT	GGG	GAC	CCT	1860
Tyr	Lys	Pro	Phe	Gln	Asp	Leu	Ile	Pro	Gly	Cys	Pro	Gly	Asp	Pro	620
GCC	CCA	GTC	CCT	GTC	CCC	TTG	TTC	ACC	TTT	GGA	CTG	GAC	AGG	GAG	1905
Ala	Pro	Val	Pro	Val	Pro	Leu	Phe	Thr	Phe	Gly	Leu	Asp	Arg	Glu	635
CCA	CCT	CGC	AGT	CCG	CAG	AGC	TCA	CAT	CTC	CCA	AGC	AGC	TCC	CCA	1950
Pro	Pro	Arg	Ser	Pro	Gln	Ser	Ser	His	Leu	Pro	Ser	Ser	Ser	Pro	650
GAG	CAC	CTG	GGT	CTG	GAG	CCG	GGG	GAA	AAG	GTA	GAG	GAC	ATG	CCA	1995
Glu	His	Leu	Gly	Leu	Glu	Pro	Gly	Glu	Lys	Val	Glu	Asp	Met	Pro	665
AAG	CCC	CCA	CTT	CCC	CAG	GAG	CAG	GCC	ACA	GAC	CCC	CTT	GTG	GAC	2040
Lys	Pro	Pro	Leu	Pro	Gln	Glu	Gln	Ala	Thr	Asp	Pro	Leu	Val	Asp	680
AGC	CTG	GGC	AGT	GGC	ATT	GTC	TAC	TCA	GCC	CTT	ACC	TGC	CAC	CTG	2085
Ser	Leu	Gly	Ser	Gly	Ile	Val	Tyr	Ser	Ala	Leu	Thr	Cys	His	Leu	695
TGC	GGC	CAC	CTG	AAA	CAG	TGT	CAT	GGC	CAG	GAG	GAT	GGT	GGC	CAG	2130
Cys	Gly	His	Leu	Lys	Gln	Cys	His	Gly	Gln	Glu	Asp	Gly	Gly	Gln	710
ACC	CCT	GTC	ATG	GCC	AGT	CCT	TGC	TGT	GGC	TGC	TGC	TGT	GGA	GAC	2175
Thr	Pro	Val	Met	Ala	Ser	Pro	Cys	Cys	Gly	Cys	Cys	Cys	Gly	Asp	725
AGG	TCC	TCG	CCC	CCT	ACA	ACC	CCC	CTG	AGG	GCC	CCA	GAC	CCC	TCT	2220
Arg	Ser	Ser	Pro	Pro	Thr	Thr	Pro	Leu	Arg	Ala	Pro	Asp	Pro	Ser	740
CCA	GGT	GGG	GTT	CCA	CTG	GAG	GCC	AGT	CTG	TGT	CCG	GCC	TCC	CTG	2265
Pro	Gly	Gly	Val	Pro	Leu	Glu	Ala	Ser	Leu	Cys	Pro	Ala	Ser	Leu	755
GCA	CCC	TCG	GGC	ATC	TCA	GAG	AAG	AGT	AAA	TCC	TCA	TCA	TCC	TTC	2310
Ala	Pro	Ser	Gly	Ile	Ser	Glu	Lys	Ser	Lys	Ser	Ser	Ser	Ser	Phe	770
CAT	CCT	GCC	CCT	GGC	AAT	GCT	CAG	AGC	TCA	AGC	CAG	ACC	CCC	AAA	2355
His	Pro	Ala	Pro	Gly	Asn	Ala	Gln	Ser	Ser	Ser	Gln	Thr	Pro	Lys	785
ATC	GTG	AAC	TTT	GTC	TCC	GTG	GGA	CCC	ACA	TAC	ATG	AGG	GTC	TCT	2400
Ile	Val	Asn	Phe	Val	Ser	Val	Gly	Pro	Thr	Tyr	Met	Arg	Val	Ser	800

Title: ANTIBODIES THAT BIND INTERLEUKIN-4
RECEPTOR

Filed: November 7, 2003

Inventors: Hongxing Zhou et al.

Docket No.: 3492

Figure 2A

FR1									
L1	1	GAAATTGTGT	TGACGCAGTC	TCCAGGCACC	CTGTCTTTGT	CTCCAGGGGA	AAGAGCCACC	60	
L2	1	GAAATTGTGT	TGACGCAGTC	TCCAGGCACC	CTGTCTTTGT	CTCCAGGGGA	AAGAGCCACC	60	
L3	1	GAAATTGTGT	TGACGCAGTC	TCCAGGCACC	CTGTCTTTGT	CTCCGGGGGA	AAGAGCCACC	60	
L4	1	GAAATTGTGA	TGACGCAGTC	TCCAGGCACC	CTGTCTTTGT	CTCCAGGGGA	AAGAGCCACC	60	
L5	1	GATATTGTGC	TGACCCAGTC	TCCAGCCACC	CTGTCTTTGT	CTCCAGGGGA	AAGAGCCACC	60	
L6	1	GATATTGTGC	TGACGCAGAC	TCCAGCCACC	CTGTCTTTGT	CTCCAGGGGA	AAGAGCCACC	60	
CDR1									
L1	61	CTCTCCTGCA	GGGCCAGTCA	GAGTGTTAGC	AGCAGCTACT	TAGCCTGGTA	CCAGCAGAAA	120	
L2	61	CTCTCCTGCA	GGGCCAGTCA	GAGTGTTAGC	AACAGCTACT	TAGCCTGGTA	CCAGCAGAAA	120	
L3	61	CTCTCCTGCA	GGGCCAGTCA	GACTGTTAAC	AGCGACTACT	TAGCCTGGTA	CCAGCAGAAA	120	
L4	61	CTCTCCTGCA	GGGCCAGTCA	GAGTGTTAGC	AGCGACTACT	TAGCCTGGTA	CCAGCAGAAA	120	
L5	61	CTCTCCTGCA	GGGCCAGTCA	GAGTGTTAAC	AGCAACTACT	TAGCCTGGTA	CCAGCAGAAA	120	
L6	61	CTCTCCTGCA	GGGCCAGTCA	GAGTGTTGGC	AGCAGCTACT	TAGCCTGGTA	CCAGCAGAGA	120	
FR2					CDR2				
L1	121	CCTGGCCAGG	CTCCCAGGCT	CCTCATCTTT	GGTGCATCCA	GCAGGGCCAC	TGGCATCCCA	180	
L2	121	CCTGGCCAGG	CTCCCAGGCT	CCTCATCTAT	GGTGCATCCA	GCAGGGCCCC	TGGCATCCCA	180	
L3	121	CCGGGCCAGG	CTCCCAGGCT	CCTCATCTAT	GGTGCATCCA	GCAGGGCCAC	TGGCATCCCA	180	
L4	121	CCTGGCCAGG	CTCCCAGGCT	CCTCATCTAT	GGTGCATCTA	GCAGGGCCTC	TGGCATCCCA	180	
L5	121	CCTGGCCAGG	CTCCCAGGCT	CCTCATCTAT	GGTACATCCT	ACAGGGCCAC	TGGCATCCCA	180	
L6	121	CCTGGCCAGG	CTCCCAGGCT	CCTCATCTAT	GGTGCATCCA	GCAGGGCCAC	TGGCATCCCG	180	
FR3									
L1	181	GACAGGTTCA	GTGGCAGTGG	GTCTGGGACA	GACTTCACTC	TCACCATCAG	CAGACTGGAG	240	
L2	181	GACAGGTTCA	GTGGCAGTGG	GTCTGGGACA	GACTTCACTC	TCACCATCAG	CAGACTGGAG	240	
L3	181	GACAGGTTCA	GTGGCAGTGG	GTCTGGGACA	GACTTCACTC	TCACCATCAG	CAGACTGGAG	240	
L4	181	GACAGGTTCA	GTGGCAGTGG	GTTTGGGACA	GACTTCACTC	TCACCATCAG	CAGACTGGAG	240	
L5	181	GACAGGTTCA	GTGGCAGTGG	GTCTGGGACA	GACTTCACTC	TCACCATCAC	CAGACTGGAG	240	
L6	181	GACAGGTTCA	GTGGCAGTGG	GTCTGGGACA	GACTTCACTC	TCACGATCAG	CAGACTGGAG	240	
CDR3									
L1	241	CCTGAAGATT	TTGCAGTGTA	TTACTGTCAG	CAGTATGGTA	GCTCACCTCC	GTGGACGTTT	300	
L2	241	CCTGAAGATT	TTGCAGTGTA	TTACTGTCAG	CAGTATGATC	ACTCAGCAGG	GTGGACGTTT	300	
L3	241	CCTGAAGATT	TTGCAGTCTA	TTACTGTCAG	CAGTATGGTA	GGTCACCTCC	GTGGACGTTT	300	
L4	241	CCTGAAGATT	TTGCAATATA	TTACTGTCAG	CAGTATGGTA	GCTCACCTCC	GTGGACGTTT	300	
L5	241	CCTGAAGATT	TTGCAGTGTA	TTACTGTCAG	CAGTATGGTA	GCTCACCACC	GTGGACGTTT	300	
L6	241	CCTGAAGATT	TTGCAGTGTA	TTATTGTCAG	CAGTATGGAA	GTTACCTCC	GTGGATGTTT	300	
FR4									
L1	301	GGCCAAGGGA	CCAAGGTGGA	AATCAAA	327				
L2	301	GGCCAAGGGA	CCAAGGTGGA	GATCAAA	327				
L3	301	GGCCAAGGGA	CCAAGGTGGA	TATCAAA	327				
L4	301	GGCCAAGGGA	CCAAGGTGGA	AATCAAA	327				
L5	301	GGCCAAGGGA	CACGACTGGA	GATTAAA	327				
L6	301	GGCCAAGGGA	CCAAGGTGGA	GATCAAA	327				

Title: ANTIBODIES THAT BIND INTERLEUKIN-4
RECEPTOR

Filed: November 7, 2003

Inventors: Hongxing Zhou et al.

Docket No.: 3492

Figure 2B

FR1									
H1	1	GAGGTTTCAGC	TGGTGCAGTC	TGGGGGAGGC	TTGGTACATC	CTGGGGGGGTC	CCTGAGACTC	60	
H2	1	GAGGTTTCAGC	TGGTGCAGTC	TGGGGGAGGC	TTGGTACATC	CTGGGGGGGTC	CCTGAGACTC	60	
H3	1	GAGGTTTCAGC	TGGTGCAGTC	TGGGGGAGGC	TTGGTACATC	CTGGGGGGGTC	CCTGAGACTC	60	
H4	1	GAGGTTTCAGT	TGGTGCAGTC	TGGGGGAGGC	TTGGTACATC	CTGGGGGGGTC	CCTGAGACTC	60	
H5	1	GAGGTTTCAGC	TGGTGCAGTC	TGGGGGAGGC	TTGGTACATC	CTGGGGGGGTC	CCTGAGACTC	60	
H6	1	GAGGTTTCAGT	TGGTGCAGTC	TGGGGGAGGC	TTGGTACATC	CTGGGGGGGTC	CCTGAGACTC	60	
H7	1	GAGGTTTCAGC	TGGTGCAGTC	TGGGGGAGGC	TTGGTACATC	CTGGGGGGGTC	CCTGAGACTC	60	
H8	1	GAGGTTTCAGT	TGGTGCAGTC	TGGGGGAGGC	TTGGTACATC	CTGGGGGGGTC	CCTGAGACTC	60	
H9	1	GAGGTTTCAGC	TGGTGCAGTC	TGGGGGAGGC	TTGGTACATC	CTGGGGGGGTC	CCTGAGACTC	60	
H10	1	GAGGTTTCAGT	TGGTGCAGTC	TGGGGGAGGC	TTGGTACATC	CTGGGGGGGTC	CCTGAGACTC	60	
H11	1	GAGGTTTCAGC	TGGTGCAGTC	TGGGGGAGGC	TTGGTACATC	CTGGGGGGGTC	CCTGAGACTC	60	
H12	1	GAGGTTTCAGC	TGGTGCAGTC	TGGGGGAGGC	TTGGTACATC	CTGGGGGGGTC	CCTGAGACTC	60	
H13	1	GAGGTTTCAGC	TGGTGCAGTC	TGGGGGAGGC	TTGGTACATC	CTGGGGGGGTC	CCTGAGACTC	60	
H14	1	GAGGTTTCAGT	TGGTGCAGTC	TGGGGGAGGC	TTGGTACAGC	CTGGGGGGGTC	CCTGAGACTC	60	
H15	1	GAGGTTTCAGT	TGGTGGAGTC	TGGGGGAGGC	TTGGTACAGC	CTGGGGGGGTC	CCTGAGACTC	60	
H16	1	GAGGTTTCAGT	TGGTGGAGTC	TGGGGGAGGC	TTGGTACAGC	CTGGGGGGGTC	CCTGAGACTC	60	
H17	1	GAGGTTTCAGT	TGGTGGAGTC	TGGGGGAGGC	TTGGTACAGC	CTGGGGGGGTC	CCTGAGACTC	60	
H18	1	GAGGTTTCAGT	TGGTGGAGTC	TGGGGGAGGC	TTGGTACAGC	CTGGGGGGGTC	CCTGAGACTC	60	
H19	1	GAGGTTTCAGT	TGGTGGAGTC	TGGGGGAGGC	TTGGTACAGC	CTGGGGGGGTC	CCTGAGACTC	60	
H20	1	GAGGTTTCAGT	TGGTGGAGTC	TGGGGGAGGC	TTGGTACAGC	CTGGGGGGGTC	CCTGAGACTC	60	
H21	1	GAGGTTTCAGT	TGGTGGAGTC	TGGGGGAGGC	TTGGTACAGC	CTGGGGGGGTC	CCTGAGACTC	60	
H22	1	GAGGTTTCAGT	TGGTGGAGTC	TGGGGGAGGC	TTGGTACAGC	CTGGGGGGGTC	CCTGAGACTC	60	
H23	1	GAGGTTTCAGT	TGGTGGAGTC	TGGGGGAGGC	TTGGTACAGC	CTGGGGGGGTC	CCTGAGACTC	60	
H24	1	GAGGTTTCAGT	TGGTGGAGTC	TGGGGGAGGC	TTGGTACAGC	CTGGGGGGGTC	CCTGAGACTC	60	

CDR1									
H1	61	TCCTGTGCAG	GCTCTGGATT	CACCTTCAGT	<u>AGAAATGCTA</u>	<u>TGTTCTGGGT</u>	TCGCCAGGCT	120	
H2	61	TCCTGTGCAG	GCTCTGGATT	CACCTTCAGT	<u>AGAAATGCTA</u>	<u>TGTTCTGGGT</u>	TCGCCAGGCT	120	
H3	61	TCCTGTGCAG	GCTCTGGATT	CACCTTCAGT	<u>AGAAATGCTA</u>	<u>TGTTCTGGGT</u>	TCGCCAGGCT	120	
H4	61	TCCTGTGCAG	GCTCTGGATT	CACCTTCAGT	<u>AGAAATGCTA</u>	<u>TGTTCTGGGT</u>	TCGCCAGGCT	120	
H5	61	TCCTGTGCAG	GCTCTGGATT	CACCTTCAGT	<u>AGAAATGCTA</u>	<u>TGTTCTGGGT</u>	TCGCCAGGCT	120	
H6	61	TCCTGTGCAG	GCTCTGGATT	CACCTTCAGT	<u>AGAAATGCTA</u>	<u>TGTTCTGGGT</u>	TCGCCAGGCT	120	
H7	61	TCCTGTGCAG	GCTCTGGATT	CACCTTCAGT	<u>AGAAATGCTA</u>	<u>TGTTCTGGGT</u>	TCGCCAGGCT	120	
H8	61	TCCTGTGCAG	GCTCTGGATT	CACCTTCAGT	<u>AGAAATGCTA</u>	<u>TGTTCTGGGT</u>	TCGCCAGGCT	120	
H9	61	TCCTGTGCAG	GCTCTGGATT	CACCTTCAGT	<u>AGAAATGCTA</u>	<u>TGTTCTGGGT</u>	TCGCCAGGCT	120	
H10	61	TCCTGTGCAG	GCTCTGGATT	CACCTTCAGT	<u>AGAAATGCTA</u>	<u>TGTTCTGGGT</u>	TCGCCAGGCT	120	
H11	61	TCCTGTGCAG	GCTCTGGATT	CACCTTCAGT	<u>AGAAATGCTA</u>	<u>TGTTCTGGGT</u>	TCGCCAGGCT	120	
H12	61	TCCTGTGCAG	GCTCTGGATT	CACCTTCAGT	<u>AGAAATGCTA</u>	<u>TGTTCTGGGT</u>	TCGCCAGGCT	120	
H13	61	TCCTGTGCAG	GCTCTGGATT	CACCTTCAGT	<u>AGAAATGCTA</u>	<u>TGTTCTGGGT</u>	TCGCCAGGCT	120	
H14	61	TCCTGTGCAG	CCTCTGGATT	CACCTTCAGT	<u>AGAAATGCTA</u>	<u>TGTTCTGGGT</u>	TCGCCAGGCT	120	
H15	61	TCCTGTGCAG	CCTCTGGATT	CACCTTCAGT	<u>AGAAATGCTA</u>	<u>TGTTCTGGGT</u>	TCGCCAGGCT	120	
H16	61	TCCTGTGCAG	CCTCTGGATT	CACCTTCAGT	<u>AGAAATGCTA</u>	<u>TGTTCTGGGT</u>	TCGCCAGGCT	120	
H17	61	TCCTGTGCAG	CCTCTGGATT	CACCTTCAGT	<u>AGAAATGCTA</u>	<u>TGTTCTGGGT</u>	TCGCCAGGCT	120	
H18	61	TCCTGTGCAG	CCTCTGGATT	CACCTTCAGT	<u>AGAAATGCTA</u>	<u>TGTTCTGGGT</u>	TCGCCAGGCT	120	
H19	61	TCCTGTGCAG	CCTCTGGATT	CACCTTCAGT	<u>AGAAATGCTA</u>	<u>TGTTCTGGGT</u>	TCGCCAGGCT	120	
H20	61	TCCTGTGCAG	CCTCTGGATT	CACCTTCAGT	<u>AGAAATGCTA</u>	<u>TGTTCTGGGT</u>	TCGCCAGGCT	120	
H21	61	TCCTGTGCAG	CCTCTGGATT	CACCTTCAGT	<u>AGAAATGCTA</u>	<u>TGTTCTGGGT</u>	TCGCCAGGCT	120	
H22	61	TCCTGTGCAG	CCTCTGGATT	CACCTTCAGT	<u>AGAAATGCTA</u>	<u>TGTTCTGGGT</u>	TCGCCAGGCT	120	
H23	61	TCCTGTGCAG	CCTCTGGATT	CACCTTCAGT	<u>AGAAATGCTA</u>	<u>TGTTCTGGGT</u>	TCGCCAGGCT	120	
H24	61	TCCTGTGCAG	CCTCTGGATT	CACCTTCAGT	<u>AGAAATGCTA</u>	<u>TGTTCTGGGT</u>	TCGCCAGGCT	120	

Title: ANTIBODIES THAT BIND INTERLEUKIN-4
RECEPTOR

Filed: November 7, 2003

Inventors: Hongxing Zhou et al.

Docket No.: 3492

Figure 2C

		FR2			CDR2		
H1	121	CCAGGAAAAG	GTCTGGAGTG	GGTATCAGGT	ATTGGTACTG	GTGGTGCCAC	AAACTATGCA 180
H2	121	CCAGGAAAAG	GTCTGGAGTG	GGTATCAGGT	ATTGGTACTG	GTGGTGCCAC	AAGCTATGCA 180
H3	121	CCAGGAAAAG	GTCTGGAGTG	GGTATCAGGT	ATTGGTACTG	GTGGTGCCAC	AAGCTATGCA 180
H4	121	CCAGGAAAAG	GTCTGGAGTG	GGTATCAGGT	ATTGGTACTG	GTGGTGCCAC	AAACTATGCA 180
H5	121	CCAGGAAAAG	GTCTGGAGTG	GGTATCAGGT	ATTGGTACTG	GTGGTGCCAC	AAACTATGCA 180
H6	121	CCAGGAAAAG	GTCTGGAGTG	GGTATCAGGT	ATTGGTACTG	GTGGTGCCAC	AAACTATGCA 180
H7	121	CCAGGAAAAG	GTCTGGAGTG	GGTATCAGGT	ATTGGTACTG	GTGGTGCCAC	AAGCTATGCA 180
H8	121	CCAGGAAAAG	GTCTGGAGTG	GGTATCAGGT	ATTGGTACTG	GTGGTGCCAC	AAACTATGCA 180
H9	121	CCAGGAAAAG	GTCTGGAGTG	GGTATCAGGT	ATTGGTACTG	GTGGTGCCAC	AAGCTATGCA 180
H10	121	CCAGGAAAAG	GTCTGGAGTG	GGTATCAGGT	ATTGGTACTG	GTGGTGCCAC	AAACTATGCA 180
H11	121	CCAGGAAAAG	GTCTGGAGTG	GGTATCAGGT	ATTGGTACTG	GTGGTGCCAC	AAGCTATGCA 180
H12	121	CCAGGAAAAG	GTCTGGAGTG	GGTATCAGGT	ATTGGTACTG	GTGGTGCCAC	AAGCTATGCA 180
H13	121	CCAGGAAAAG	GTCTGGAGTG	GGTATCAGGT	ATTGGTACTG	GTGGTGCCAC	AAGCTATGCA 180
H14	121	CCAGGAAAAG	GTCTGGAGTG	GGTATCAGGT	ATTGGTACTG	GTGGTGCCAC	AAGCTATGCA 180
H15	121	CCAGGAAAAG	GTCTGGAGTG	GGTATCAGGT	ATTGGTACTG	GTGGTGCCAC	AAGCTATGCA 180
H16	121	CCAGGAAAAG	GTCTGGAGTG	GGTATCAGGT	ATTGGTACTG	GTGGTGCCAC	AAGCTATGCA 180
H17	121	CCAGGAAAAG	GTCTGGAGTG	GGTATCAGGT	ATTGGTACTG	GTGGTGCCAC	AAGCTATGCA 180
H18	121	CCAGGAAAAG	GTCTGGAGTG	GGTATCAGGT	ATTGGTACTG	GTGGTGCCAC	AAGCTATGCA 180
H19	121	CCAGGAAAAG	GTCTGGAGTG	GGTATCAGGT	ATTGGTACTG	GTGGTGCCAC	AAGCTATGCA 180
H20	121	CCAGGAAAAG	GTCTGGAGTG	GGTATCAGGT	ATTGGTACTG	GTGGTGCCAC	AAGCTATGCA 180
H21	121	CCAGGAAAAG	GTCTGGAGTG	GGTATCAGGT	ATTGGTACTG	GTGGTGCCAC	AAGCTATGCA 180
H22	121	CCAGGAAAAG	GTCTGGAGTG	GGTATCAGGT	ATTGGTACTG	GTGGTGCCAC	AAGCTATGCA 180
H23	121	CCAGGAAAAG	GTCTGGAGTG	GGTATCAGGT	ATTGGTACTG	GTGGTGCCAC	AAGCTATGCA 180
H24	121	CCAGGAAAAG	GTCTGGAGTG	GGTATCAGGT	ATTGGTACTG	GTGGTGCCAC	AAGCTATGCA 180

					FR3		
H1	181	GACTCCGTGA	AGGGCCGATT	CACCATCTCC	AGAGACAATG	CCAAGAACTC	CTTGTATCTT 240
H2	181	GACTCCGTGA	AGGGCCGATT	CACCATCTCC	AGAGACAATG	CCAAGAACTC	CTTGTATCTT 240
H3	181	GACTCCGTGA	AGGGCCGATT	CACCATCTCC	AGAGACAATG	CCAAGAACTC	CTTGTATCTT 240
H4	181	GACTCCGTGA	AGGGCCGATT	CACCATCTCC	AGAGACAATG	CCAAGAACTC	CTTGTATCTT 240
H5	181	GACTCCGTGA	AGGGCCGATT	CACCATCTCC	AGAGACAATG	CCAAGAACTC	CTTGTATCTT 240
H6	181	GACTCCGTGA	AGGGCCGATT	CACCATCTCC	AGAGACAATG	CCAAGAACTC	CTTGTATCTT 240
H7	181	GACTCCGTGA	AGGGCCGATT	CACCATCTCC	AGAGACAATG	CCAAGAACTC	CTTGTATCTT 240
H8	181	GACTCCGTGA	AGGGCCGATT	CACCATCTCC	AGAGACAATG	CCAAGAACTC	CTTGTATCTT 240
H9	181	GACTCCGTGA	AGGGCCGATT	CACCATCTCC	AGAGACAATG	CCAAGAACTC	CTTGTATCTT 240
H10	181	GACTCCGTGA	AGGGCCGATT	CACCATCTCC	AGAGACAATG	CCAAGAACTC	CTTGTATCTT 240
H11	181	GACTCCGTGA	AGGGCCGATT	CACCATCTCC	AGAGACAATG	CCAAGAACTC	CTTGTATCTT 240
H12	181	GACTCCGTGA	AGGGCCGATT	CACCATCTCC	AGAGACAATG	CCAAGAACTC	CTTGTATCTT 240
H13	181	GACTCCGTGA	AGGGCCGATT	CACCATCTCC	AGAGACAATG	CCAAGAACTC	CTTGTATCTT 240
H14	181	GACTCCGTGA	AGGGCCGATT	CACCATCTCC	AGAGACAATG	CCAAGAACTC	CTTGTATCTT 240
H15	181	GACTCCGTGA	AGGGCCGATT	CACCATCTCC	AGAGACAATG	CCAAGAACTC	CTTGTATCTT 240
H16	181	GACTCCGTGA	AGGGCCGATT	CACCATCTCC	AGAGACAATG	CCAAGAACTC	CTTGTATCTT 240
H17	181	GACTCCGTGA	AGGGCCGATT	CACCATCTCC	AGAGACAATG	CCAAGAACTC	CTTGTATCTT 240
H18	181	GACTCCGTGA	AGGGCCGATT	CACCATCTCC	AGAGACAATG	CCAAGAACTC	CTTGTATCTT 240
H19	181	GACTCCGTGA	AGGGCCGATT	CACCATCTCC	AGAGACAATG	CCAAGAACTC	CTTGTATCTT 240
H20	181	GACTCCGTGA	AGGGCCGATT	CACCATCTCC	AGAGACAATG	CCAAGAACTC	CTTGTATCTT 240
H21	181	GACTCCGTGA	AGGGCCGATT	CACCATCTCC	AGAGACAATG	CCAAGAACTC	CTTGTATCTT 240
H22	181	GACTCCGTGA	AGGGCCGATT	CACCATCTCC	AGAGACAATG	CCAAGAACTC	CTTGTATCTT 240
H23	181	GACTCCGTGA	AGGGCCGATT	CACCATCTCC	AGAGACAATG	CCAAGAACTC	CTTGTATCTT 240
H24	181	GACTCCGTGA	AGGGCCGATT	CACCATCTCC	AGAGACAATG	CCAAGAACTC	CTTGTATCTT 240

Title: ANTIBODIES THAT BIND INTERLEUKIN-4
RECEPTOR

Filed: November 7, 2003

Inventors: Hongxing Zhou et al.

Docket No.: 3492

Figure 2D

H1	241	CAAATGAACA	GCCTGAGAGC	CGAGGACATG	GCTGTGTATT	ACTGTGCAAG	AGGGAGGTAC	300
H2	241	CAAATGAACA	GCCTGAGTGC	CGAGGACATG	GCTGTGTATT	ACTGTGCAAG	AGGGAGGTAC	300
H3	241	CAAATGAACA	GCCTGAGAGC	CGAGGACATG	GCTGTGTATT	ACTGTGCAAG	AGGGAGGTAC	300
H4	241	CAAATGAACA	GCCTGAGAGC	CGAGGACATG	GCTGTGTATT	ACTGTGCAAG	AGGGAGGTAC	300
H5	241	CAAATGAACA	GCCTGAGAGC	CGAGGACATG	GCTGTGTATT	ACTGTGCAAG	AGGGAGGTAC	300
H6	241	CAAATGAACA	GCCTGAGAGC	CGAGGACATG	GCTGTGTATT	ACTGTGCAAG	AGGGAGGTAC	300
H7	241	CAAATGAACA	GCCTGAGAGC	CGAGGACATG	GCTGTGTATT	ACTGTGCAAG	AGGGAGGTAC	300
H8	241	CAAATGAACA	GCCTGAGAGC	CGAGGACATG	GCTGTGTATT	ACTGTGCAAG	AGGGAGGTAC	300
H9	241	CAAATGAACA	GCCTGAGAGC	CGAGGACATG	GCTGTGTATT	ACTGTGCAAG	AGGGAGGTAC	300
H10	241	CAAATGAACA	GCCTGAGAGC	CGAGGACATG	GCTGTGTATT	ACTGTGCAAG	AGGGAGGTAC	300
H11	241	CAAATGAACA	GCCTGAGAGC	CGAGGACATG	GCTGTGTATT	ACTGTGCAAG	AGGGAGGTAC	300
H12	241	CAAATGAACA	GCCTGAGAGC	CGAGGACATG	GCTGTGTATT	ACTGTGCAAG	AGGGAGGTAC	300
H13	241	CAAATGAACA	GCCTGAGTGC	CGAGGACATG	GCTGTGTATT	ACTGTGCAAG	AGGGAGGTAC	300
H14	241	CAAATGAACA	GCCTGAGAGC	CGAGGACACG	GCTGTGTATT	ACTGTGCAAG	AGGGAGGTAC	300
H15	241	CAAATGAACA	GCCTGAGAGC	CGAGGACACG	GCTGTGTATT	ACTGTGCAAG	AGGGAGGTAC	300
H16	241	CAAATGAACA	GCCTGAGAGC	CGAGGACACG	GCTGTGTATT	ACTGTGCAAG	AGGGAGGTAC	300
H17	241	CAAATGAACA	GCCTGAGAGC	CGAGGACACG	GCTGTGTATT	ACTGTGCAAG	AGGGAGGTAC	300
H18	241	CAAATGAACA	GCCTGAGAGC	CGAGGACACG	GCTGTGTATT	ACTGTGCAAG	AGGGAGGTAC	300
H19	241	CAAATGAACA	GCCTGAGAGC	CGAGGACACG	GCTGTGTATT	ACTGTGCAAG	AGGGAGGTAC	300
H20	241	CAAATGAACA	GCCTGAGAGC	CGAGGACACG	GCTGTGTATT	ACTGTGCAAG	AGGGAGGTAC	300
H21	241	CAAATGAACA	GCCTGAGAGC	CGAGGACACG	GCTGTGTATT	ACTGTGCAAG	AGGGAGGTAC	300
H22	241	CAAATGAACA	GCCTGAGAGC	CGAGGACACG	GCTGTGTATT	ACTGTGCAAG	AGGGAGGTAC	300
H23	241	CAAATGAACA	GCCTGAGAGC	CGAGGACACG	GCTGTGTATT	ACTGTGCAAG	AGGGAGGTAC	300
H24	241	CAAATGAACA	GCCTGAGAGC	CGAGGACACG	GCTGTGTATT	ACTGTGCAAG	AGGGAGGTAC	300

		CDR3		FR4		
H1	301	TACTTTGACT	ACTGGGGCCA	GGGAACCCCTG	GTCACCGTCT	CCTCA 345
H2	301	TACTTCACCC	ACTGGGGCCA	GGGAACCCCTG	GTCACCGTCT	CCTCA 345
H3	301	TGGTACAACA	ACTGGGGCCA	GGGAACCCCTG	GTCACCGTCT	CCTCA 345
H4	301	TACTTCCCGT	GGTGGGGCCA	GGGAACCCCTG	GTCACCGTCT	CCTCA 345
H5	301	TACTTCACGA	GGTGGGGCCA	GGGAACCCCTG	GTCACCGTCT	CCTCA 345
H6	301	TGGTACCCGT	GGTGGGGCCA	GGGAACCCCTG	GTCACCGTCT	CCTCA 345
H7	301	TGGTACCCGT	GGTGGGGCCA	GGGAACCCCTG	GTCACCGTCT	CCTCA 345
H8	301	TGGTTCCCGT	GGTGGGGCCA	GGGAACCCCTG	GTCACCGTCT	CCTCA 345
H9	301	TGGTTCCCGT	GGTGGGGCCA	GGGAACCCCTG	GTCACCGTCT	CCTCA 345
H10	301	TGGTACCCGT	GGTGGGGCCA	GGGAACCCCTG	GTCACCGTCT	CCTCA 345
H11	301	TGGTACCCGT	GGTGGGGCCA	GGGAACCCCTG	GTCACCGTCT	CCTCA 345
H12	301	TACTTCCCGT	GGTGGGGCCA	GGGAACCCCTG	GTCACCGTCT	CCTCA 345
H13	301	TACTTCCCGT	GGTGGGGCCA	GGGAACCCCTG	GTCACCGTCT	CCTCA 345
H14	301	TACTTCCCGT	GGTGGGGCCA	GGGAACCCCTG	GTCACCGTCT	CCTCA 345
H15	301	TACTTTGACT	ACTGGGGCCA	GGGAACCCCTG	GTCACCGTCT	CCTCA 345
H16	301	TACTTCACCC	ACTGGGGCCA	GGGAACCCCTG	GTCACCGTCT	CCTCA 345
H17	301	TGGTACAACA	ACTGGGGCCA	GGGAACCCCTG	GTCACCGTCT	CCTCA 345
H18	301	TACTTCACGA	GGTGGGGCCA	GGGAACCCCTG	GTCACCGTCT	CCTCA 345
H19	301	TGGTACCCGT	GGTGGGGCCA	GGGAACCCCTG	GTCACCGTCT	CCTCA 345
H20	301	TGGTACCCGT	GGTGGGGCCA	GGGAACCCCTG	GTCACCGTCT	CCTCA 345
H21	301	TGGTTCCCGT	GGTGGGGCCA	GGGAACCCCTG	GTCACCGTCT	CCTCA 345
H22	301	TGGTTCCCGT	GGTGGGGCCA	GGGAACCCCTG	GTCACCGTCT	CCTCA 345
H23	301	TGGTACCCGT	GGTGGGGCCA	GGGAACCCCTG	GTCACCGTCT	CCTCA 345
H24	301	TGGTACCCGT	GGTGGGGCCA	GGGAACCCCTG	GTCACCGTCT	CCTCA 345

Title: ANTIBODIES THAT BIND INTERLEUKIN-4
RECEPTOR

Filed: November 7, 2003

Inventors: Hongxing Zhou et al.

Docket No.: 3492

Figure 3

	FR1	CDR1	FR2	CDR2	FR3	CDR3	FR4	
L1	EIVLTQSPGTL	SLSPGERATL	SCRASQSVSS	YLAWYQOKPGQAPRL	IFGASSRATG	IPDRFSGSGGTDF	TLISRLEPEDFAVYVCQQYGS	SPFTFGCGTKVEIK
L2	---	N	---	Y	---	P	---	DH-AG
L3	---	T-N-D	---	Y	---	Y	---	R
L4	M	D	---	Y	---	S	---	F
L5	D	N-N	---	Y-T-Y	---	T	---	RL
L6	D	T-A	---	R	---	Y	---	M
H1	EVQLVQSGGGLVHPGGS	LRLSCAGSGFTFS	RNAMPVWRQAPGKGL	EWYSGIGTGGATYADSV	TKGRFTISRDNAKNSLYLQWNS	LRAEDMAVYVCARGY	YFDYWGQGLVT	VSS
H2	---	---	---	S	---	S	---	TH
H3	---	---	---	S	---	WTNN	---	FW
H4	---	---	---	---	---	FW	---	TR
H5	---	---	---	---	---	YFW	---	YFW
H6	---	---	---	S	---	YFW	---	W-FW
H7	---	---	---	S	---	W-FW	---	W-FW
H8	---	---	---	S	---	W-FW	---	W-FW
H9	---	---	---	S	---	W-FW	---	W-FW
H10	---	---	---	S	---	W-FW	---	W-FW
H11	---	---	---	S	---	W-FW	---	W-FW
H12	---	---	---	S	---	W-FW	---	W-FW
H13	---	---	---	S	---	W-FW	---	W-FW
H14	E	Q	A	---	S	---	T	---
H15	E	Q	A	---	S	---	T	---
H16	E	Q	A	---	S	---	T	---
H17	E	Q	A	---	S	---	T	---
H18	E	Q	A	---	S	---	T	---
H19	E	Q	A	---	S	---	T	---
H20	E	Q	A	---	S	---	T	---
H21	E	Q	A	---	S	---	T	---
H22	E	Q	A	---	S	---	T	---
H23	E	Q	A	---	S	---	T	---
H24	E	Q	A	---	S	---	T	---

Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/US04/037242

International filing date: 04 November 2004 (04.11.2004)

Document type: Certified copy of priority document

Document details: Country/Office: US
Number: 60/518,166
Filing date: 07 November 2003 (07.11.2003)

Date of receipt at the International Bureau: 05 January 2005 (05.01.2005)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse